

IV

92
577
B82X
pt 4
Ent.

NOMENCLATURE

OF

COLEOPTEROUS . INSECTS

IN

THE COLLECTION

OF THE

BRITISH MUSEUM. ✓

Smithsonian Institution
440
Natural History Museum
P. E. Gray

SMITHSONIAN INSTITUTION

PART IV. CLERIDÆ.

8061

a. White.

PRINTED BY ORDER OF THE TRUSTEES.

LONDON: 1849.

S. J. 1913
91.745

LONDON:
SPOTTISWOODES and SHAW,
New-street-Square.

P R E F A C E .

THE object of this Nomenclature is to present a Catalogue of the well-established species of each family, with a reference to where they are figured, and a few of the more important synonyma.

It is intended for the following purposes :

1. As a List of the Species contained in the Museum collection, with an indication of their origin.
2. To point out the Desiderata of that collection at the period of publication.
3. As Labels for the Cabinet.

This list must only be considered as prefatory to a more complete Catalogue.

The family of *Cleridæ* is chiefly arranged after the method of the Marquis Spinola in his "Essai Monographique," with additions from the Monograph of Dr. Klug in the Berlin Transactions, and from other subsequent authorities.

Mr. Adam White has taken every care in the determination of the species and the references to the synonyma and figures.

JOHN EDWARD GRAY.

British Museum,
June, 1849.

THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

THE JOURNAL OF THE
THE JOURNAL OF THE
THE JOURNAL OF THE

LIST OF CLERIDÆ.

Family CLERIDÆ.

CYLIDRUS *Latr.*

- | | | |
|---|-------------------------------------|-------|
| C. CYANEUS <i>Latr.</i> | Isle of France. | B. M. |
| <i>Cuv. R. A.</i> iv. 476. | Madagascar. | |
| <i>Spin. Cler.</i> i. 85. t. 1. f. 3. | | |
| <i>Klug, Berlin. Abh.</i> 1842. p. 262. | | |
| Clerus c. <i>Fabr. Ent. Syst.</i> ii. | India (Neemuch). | |
| 209-6. | General Thomas Hardwicke. | |
| C. PALLIPES <i>Chevr.</i> | Tahiti, Java. | B. M. |
| | M. Chevrolat; Mus. "E. Ind. House." | |
| C. NIGRINUS <i>White.</i> | V. D. Land. | |
| (See Appendix.) | Coll. W. W. Saunders; F. Parry. | |
| C. BUQUETHI <i>Guér.</i> | Senegal. | |
| <i>Icon. R. A.</i> t. 15. f. 12. | | |
| <i>Spin.</i> i. 86. t. 1. f. 1. | | |
| C. BALTEATUS <i>Klug.</i> | S. Africa. | |
| <i>Berl. Abhand.</i> 1842. 263. t. 2. | | |
| f. 1. | | |
| C. FASCIATUS <i>Lap.</i> | Madagascar. | |
| <i>Silb. Rev. Ent.</i> iv. 47. | Senegal. | |
| <i>Spin.</i> i. 88. t. 1. f. 2. | | |
| <i>Klug, Berl. Abh.</i> 1842. 263. | | |
| C. succinctus <i>Dej.</i> | S. Africa. | |
| Var. C. megacephalus <i>Buq.</i> | | |
| Var. C. bimaculatus <i>Dup.</i> | | |

C. ABDOMINALIS *Klug.*

"Brasil" ?

Berl. Abhand. 1842, 262.*Cyl. fasciatus* var. *Spin.* II. 122.DENOPS *Stev.*D. PERSONATUS *Spin.*

Sicily.

B. M.

I. 90. t. 1. f. 4.

Tillus p. *Géné de quib. col.* 14. Italy.

A. Melly, Esq.

T. albofasciatus *Charp. Hor. Ent.* Banks of Rhine.

I. 198. t. 6. f. 3.

Cylidrus albof. *Klug, Berl. Abh.*

1842. 264.

T. tenuicollis *Dup.*

Crimea.

D. longicollis *Stev. Bull. Mosc.* Levant.

I. 67.

T. agilis *Lucas, Ann. Soc. Ent.* Barbary.*Fr.* 1843. 29.TILLUS *Fabr.*T. ELONGATUS *Fabr.*

Europe (Brit. Islès). B. M.

Ent. Syst. I. 78.*Oliv. t.* 1. f. 1.*Curtis, Brit. Ent.* VI. 267.*Spin. t.* 2. f. 2.*Klug, Berl. Abh.* 1842. 268.*Chrysomela elongata* *Linn. S.**Nat.* II. 603.*Lagria ruficollis* *Herbst, Fuessly,**Arch.* IV. 68. t. 23. f. 35.Var. *Spin. t.* 3. f. 1.*T. ambulans* *Fabr. Ent. Syst.* I. 78.

B. M.

Curtis, Brit. Ent. VI. 267.Var. *Spin. t.* 2. f. 3.*T. hyalinus* *Sturm, D. F. t.* 228.f. a A., *Spin. t.* 2. f. 3.*T. bimaculatus* *Donov. Brit. Ins.*

XII. t. 411. f. 2.

Cryptocephalus Marchiæ *Gmel. I.*

4. 1731.

T. UNIFASCIATUS *Latr.*

Europe (Brit. Isles). B. M.

Gen. Cr. et Ins. I. 269.*Sturm, D. F. t.* 228. f. b M.*Curtis, Brit. Ent.* VI. t. 267.*Spin. t.* 2. f. 4.*Klug, Berl. Abh.* 1842. 275.

- Cler. u. *Fabr. S. E.* i. 281.
 Tilloidea unif. *Steph. Man.* 197.
 Attelabus formicarius minor *Sulzer, Gesch. Ins.* 42. t. 4. f. 13.
 Attelab. serraticornis *De Villars, Ent.* i. 222.
 Var. T. tricolor *Dej. Spin.* t. 2. f. 5.
- T. NOTATUS *Klug.* E. Indies.
Berl. Abhand. 1842. 276.
- T. BIPARTITUS *Hombron & Jacquinet.* Guam. B. M.
Voy. au Pole Sud, t. 4. f. 13. N. Holl.
- T. CARUS. S. Australia. B. M.
 Clerus c. *Newman, Ent.* 15. Ent. Club of London.
 Opilus zonatus *Hombr. & Jacq.?*
Voy. Pole Sud, t. 4. f. 14.
- T. COLLARIS *Dej.* N. America. B. M.
Spin. 98. t. 2. f. 6. var. Columbia. W. W. Saunders, Esq.
- T. RUBRICOLLIS *Guérin.* Sennaar.
Icon. R. A. t. 15. f. 8.
Klug, Berl. Abh. 1842. 274.
Spin. t. 3. f. 3.
 T. pubescens *Spin.* i. 99. Senegal.
 Tilloidea p. *Lap. Silb. Rev. Ent.*
 Cler. uncinatus *Dup.*
 Not. Hanetii *Dej.*
- T. TRANSVERSALIS *Spin.* Coasts of Mediterranean.
 102. t. 2. f. 1. B. M.
Klug, Berl. Abh. 1842. 275. N. Africa. M. Chevrolat.
 Cler. t. *Charp. Hor. Ent.* i. t. 6. Levant. Prof. Edw. Forbes.
 f. 2. Brit. Islands.
 Cler. myrmecodes *Hoffm. Dej.*
Cat. 127.
 Cler. unifasciatus *Rossi, F. E.*
 i. 138.; *Oliv.* t. 2. f. 21.
- T. SUCCINCTUS *Dup.* E. Indies.
Spin. i. 100. t. 3. f. 4.
- T. COMPRESSICORNIS *Klug.* Cape of Good Hope.
Berl. Abh. 1842. 273. t. 2. f. 3.
- T. HILARIS *Westwood.* Australia (Sydney). B. M.
 (See Appendix.) C. Darwin, Esq.

- T. PICIPENNIS *Westwood*. E. Indies. B. M.
(See Appendix.) Col. Hearsay,
- T. BIFASCIELLUS *White*. Philippine Islands. B. M.
(See Appendix.)
- T. DISTINCTUS *White*. Java. Coll. E. Ind. House.
(See Appendix.)
- PERILYPUS *Spinola*. (*Rev. Zool. Guérin*, March, 1841.)
- P. CARBONARIUS (*Dup.*) Mexico, California.
Spin. I. 105. t. 5. f. 4.
- CALLITHERES *Dej.* (*Spinola, Rev. Zool. Guér.* March, 1841).
- C. ACUTIPENNIS *Spin.* Madagascar.
I. 108. t. 4. f. 1.
Iodamus a. Lap. Silb. Rev. Ent.
IV. 3.
Tillus a. Klug, Berl. Abh. XII. 278.
C. *Joannisii Dej.*
- C. BICOLOR *Spin.* Madagascar.
II. 127. t. 46. f. 7.
- C. TRICOLOR *Spin.* Madagascar.
I. 110. t. 3. f. 6.
Pallenis tr. Lap. Silb. Rev. Ent.
IV. 40.
Tillus tr. Klug, Berl. Abh. 1842.
276.
- C. AULICUS. Madagascar.
Tillus aul. Klug, Berl. Abhand.
1842. 277.
- C. VIDUUS. Madagascar.
Tillus vid. Klug, Berl. Abh. 1842.
278.
- C. ? 9-MACULATUS. in Gum animé.
Tillus 9-macul. Hope, Trans. Ent.
Soc. II. 54. t. 7. f. 6.
- C. LOUVELII (*Dej.*) Senegal.
Spin. I. 111. t. 3. f. 5.

PRIOCERA Kirby.

- P. VARIEGATA* Kirb. Brazil. Coll. Ent. Soc. Lond.
 Linn. Trans. XII. p. 392. t. 21. (olim. Kirby).
 f. 7. Coll. Parry.
Guérin, Icon. R. A. t. f. 10.
Spin. t. 4. f. 2.
Klug, Berl. Abh. 1842. p. 283.
- P. PUSTULATA* Spin. Mexico.
 118. t. 4. f. 4.
- P. TRINOTATA* Klug. Columbia. B. M.
Berl. Abh. 1842. p. 283.
Axina villosa Goudot. Goudot's Coll.
- P. CASTANEA.* N. Amer. B. M.
Opilus castaneus Newman, Ent. Ent. Club of London.
Mag. v. 380. Edw. Doubleday, Esq.
P. rufescens (Dej.) Spin. i. 119.
 t. 4. f. 3. ? Brazil.
Not. Leprieurii Bug.
- P. SPINOSA* Klug. Brazil. B. M.
Berl. Abh. 1842. 284. Coll. Miers.
Spin. t. 5. f. 1.
Clerus spin. Fabr. S. El. i. 280.
P. decorata Spin. 120. Brazil.
Tillus d. Dej.
T. sexpunctatus Lap. Silb. Rev.
Ent. iv. 36.
- P. MIERSIANA* White. Brazil. Coll. Miers.
 (See Appendix.)
- P. MARGINICOLLIS* Chevr. Brazil.
Thanasimus m. Chevr.
Ann. Soc. Ent. Fr. 1843.
 p. 31.
- P. CINCTIVENTRIS* Chevr. Brazil.
Thanasimus c. Chevr. Ann. Soc.
Ent. Fr. 1843. 32.
- P. ABDOMINALIS.*
Tillus abdominalis Blanchard, Bolivia (Guarayos).
D'Orbigny, Voy. à l'Amér.
Mér. vi. 91. t. 6. f. 5.

- P. BISPINOSA *Klug.* Columbia.
Berl. Abh. 1842. p. 284.
Spin. t. 18. f. 2. II. 132.
P. *Reichei Spin.* I. 121.

AXINA *Kirby.*

- A. ANALIS *Kirby.* Brazil. Coll. Ent. Soc. Lond.
Linn. Trans. XII. 391. t. 21. f. 6.
Spin. I. 123. t. 5. f. 2. (male.)
Not. crassipes *Dup.*
A. rufitarsis *Perty, Del.*
30. t. 6. f. 16. (*Notoxus ruf.*)

- A. SEX-MACULATA *Spinola.* Brazil.
I. 124. t. 5. f. 3.
Tillus sex-m. Dej.

STENOCYLIDRUS *Spinola.* (II. 129. 1844.) *Xylobius Guér. Spin.* olim, nec Latr.

- S. AZUREUS *Spin.* t. 7. f. 2. Madagascar. B. M.
Xylobius az. Spin. 128.
Tillus az. Klug, Berl. Abh.
1832. 158. t. 3. f. 6.
Var. *Tillus venustus Klug, Berl.*
Abh. 1842. 279.
♀ *Tillus longulus Klug, l. c.* 279.
Tillus pulchellus Klug, l. c. 280.

- S. ELEGANS *Spin.* Madagascar.
Xylobius el. Guér. Icon. R. A.
Spin. I. 130. t. 6. f. 1.

STENOCLERUS, n. g.

- St. FASTIGIATUS. Madagascar.
Tillus fastigiatus Klug, Berl.
Abh. 1842. 280. t. 1. f. 1.

TILLOCLERUS, n. g.

- T. AURICOMUS. Madagascar.
Tillus auricomus Klug, Berl.
Abh. 1842. 281.

PHILOCALUS *Klug.*

- P. SUCCINCTUS. C. G. Hope.
 Tillus s. *Klug, Berl. Abh.*
 1842. 281. t. 2. f. 1. (nec
 Spin.)
- P. ZONATUS. C. G. Hope.
 Tillus zon. *Klug, Berl. Abh.*
 1842. p. 282.

SYSTEMODERES *Spin. (Rev. Zool. Guér. March, 1841.)*

- S. AMENUS (*Dup.*) Mexico.
Spin. 131. t. 8. f. 1, 2.
- S. VIRIDIPENNIS (*Dup.*) Columbia.
Spin. 132. t. 8. f. 3.
- S. DASYTOIDES, n. s. Mexico. B. M.
 Clerus D. *Westw. MS.* G. R. Waterhouse, Esq.

CLERONOMUS *Klug.*

- C. BIMACULATUS. C. G. Hope.
 Tillus bim. *Klug, Berl. Abh.*
 1842. p. 281. t. 2. f. 5.

COLYPHUS *Dupont. (Rev. Zool. Guér. March, 1831.)*

- C. SIGNATICOLLIS (*Dup.*) California.
Spin. I. 135. t. 5. f. 5.
- C. CINCTIPENNIS (*Dup.*) California.
Spin. 136. t. 5. f. 6.
- C. RUFIPENNIS (*Dup.*) California.
Spin. 136. t. 9. f. 2.
- C. TERMINALIS (*Dup.*) Columbia.
Spin. 137. t. 9. f. 1.
- C. INTERCEPTUS (*Dup.*) California.
Spin. 137. t. 9. f. 3. ("inter-
 ruptus.")

PHONIUS *Chevr.*

- P. SANGUINIPENNIS *Chevr.* Mexico.
Mag. de Zool. 1843. 121. t. 107.

DERESTENUS *Chevr.*

- D. QUADRILINEATUS *Chevr.* Mexico.
Mag. de Zool. 1843. 14. t. 108.

CYMATODERA *G. R. Gray.*

- C. HOPEI *G. R. Gray.* Mexico. Coll. W. W. Saunders.
Griff. A. K. Ins. I. 375. t. 48.
 f. 1.
Lap. Silb. Rev. Zool. IV. 37.
Spin. 141. t. 9. f. 5.
Not. giganteus Dej.
Tillus Hopei Klug, Berl. Abh.
 1842. 270.
- C. IBIDIODES *Spin.* Brazil.
 II. 124. t. 47. f. 1.
- C. MARMORATA *Spin.* Mexico.
 t. 9. f. 4.
C. undata Spin. 142.
Not. und. Dup.
Not. angustatus Buq.
Not. Miletus Lap.
Tillus marmoratus Klug, Berl.
Abhand. 1842. 271.
- C. LÆTA *Spin.* Columbia. Coll. Parry.
 I. 143. t. 10. f. 4.
Not. lætus Dej.
Not. regalis Dup.
- C. PULCHELLA *White.* Columbia. B. M.
- C. PROLIXA *Spin.* Mexico. C. W. F. Parry
 Columbia.
 t. 10. f. 2. A. B.
C. modesta Spin. 144.
Not. modestus Dej. ♀
Not. terebrans Dej. ♂
Not. confusus Dup.
Tillus prolixus Klug, Berl. Abh.
 1842. 272.
- C. CONFLAGRATA. S. America (Mission of Ori-
Tillus confl. Klug, Berlin Abh. noco).
 1842.

- C. LONGICOLLIS *Spin.* N. America.
 147. t. 10. f. 1.
 Not. l. *Dej.*
 Var. Not. Vauthieri *Dup.*
 Var. Not. brunneus *Dej.*
- C. CYLINDRICOLLIS *Chevr.* Mexico.
Col. Mex. fasc. i.
Spin. i. 148. t. 10. f. 3.
 Not. cyl. *Lap.*
 Not. brunneus *Dej.*
 Tillus cyl. *Klug, Berl. Abh.* 1842.
 271.
- C. INORNATA. N. America. B. M.
Priocera in. Say, Boston Journ. Ent. Club of London.
Nat. Hist. i. 161.
 Tillus in. *Klug, Berl. Abh.* 1842.
 271.
- C. BOSCH *Chevr.* N. America.
Ann. Soc. Ent. France, 1843.
 p. 31.
- C. ANGUSTATA *Spin.* California.
 149. t. 7. f. 1.
 Not. ang. *Dej.*
- C. CINGULATA. S. Africa.
 Tillus cingulatus *Klug, Berl.*
Abh. 1842. p. 273.
- C. PALLIDIPENNIS *Chevr.* Mexico.
Mag. de Zool. 1843. 8.
- C. DISCOIDALIS *Chevr.* Mexico.
Mag. de Zool. 1843. 10.
- C. WESTWOODII *Chevr.* Mexico.

EPICLINES *Chevr.*

- E. GAYI *Chevr.* Chili (Valparaiso).
Mag. Zool. t. 231. f. 1.
Blanchard, D'Orbigny, Voy. à
l'Amér. MÉR. VI. 95. t. 6. f. 6.
 C. Gayi *Spin.* 149.
 Not. chilensis *Gory.*

- E. BASALIS *Blanch.* Chili.
D'Orbigny, Voy. à l'Amér. Mér.
 VI. 96.
 E. Gayi var?
- ELEALE *Newman.* (*Entomologist*, Jan. 1841.)
Chalciclerus Spinola.
- ELEALE ASPERA. Australia (Adelaide). B. M.
C. unicolor Spin. I. 279. t. 21. N. Holland (Van Diemen's
 f. 5.? Land. Ent. Club of London.
Eleale aspera Newman, Ent. 36.
Eleale rugosa Newman, Ent. 364.
- XYLOTRETUS *Guér. Spin.*
(Xilotretus Guér. Spin. March, 1841.)
- X. VIRIDIS *Guérin* N. Holland.
Icon. R. A. Expl. III. 52.
 (1829—1844).
Spin. 153. t. 6. f. 2.
- X. SCROBILATUS *Spin.* N. Holland.
 156.
X. foveolatus Spin. t. 15. f. 4.
- X. REICHEI *Spin.* N. Holland (Swan River).
 155. t. 7. f. 3.
X. virescens Dup.
- X. OBSCURUS *Germ.* Australia (Adelaide). B. M.
Linn. Ent. III. 185. Ent. Club of Lond.
Clerus obsc. Newman, Ent. 16.
- X. SIMPLEX.
Xanthoceros simplex Newman, Australia. B. M.
Entom. 363. Ent. Club of London.
Clerus simplex Newman, En- (Adelaide.)
tom. 16.
X. leucaspis Germ. Linn. Ent.
 III. 186.
- XYL. IDONEÛS. Australia (P. Philip). B. M.
Xanthoceros idoneus Newman, Ent. Club of London.
Entom. 363. (Aug. 1842.)

XYL. PULCHER.

Xanthoceros pulcher *Newman*,
Entom. 364.

Clerus pulcher *Newm. Ent.* 16.

Chalc. bimaculatus, *Spin.* i. 281.
 t. 20. f. 1.

Australia (Adelaide). B. M.
 Ent. Club of London.

SCROBIGER *Spinola. (Rev. Zool. Guér. March, 1841.)*
 Xanthoceros *Newman*.

S. SPLENDIDUS *Spin.*
 i. 232.

Germar, Linn. Ent. iii. 186.

Cler. splen. *Newman, Entom.* (Swan River.)
 p. 15.

Xanthoceros splen. *Newman*, (Adelaide.)
Entom. p. 363.

Scrob. Reichei *Spin.* t. 14. f. 1. W. Spry, Esq.; Ent. Club of Lond.

PTYCHOPTERUS *Klug.*

P. DIMIDIATUS *Klug.*

Berl. Abh. 1842. 316. t. 1. f. 5.

S. Africa.

TILLICERA *Spin. (Rev. Zool. Guér. March, 1841.)*

T. JAVANA *Spin.*

160. t. 12. f. 2.

Cler. jav. *Dej.*

Java.

T. CHALYBEA *Westw. n. s.*

Tennasserim. B. M.
 From Dr. Packman's Collection.

T. MUTILLÆCOLOR *White, n. s.*

N. India. B. M.
 From Capt. Boyes's Collect.

CLADISCUS *Chevr. (Ann. Soc. Ent. France, 1843. p. 33.)*

C. STRANGULATUS *Chevr.*

i. c. p. 33.

T. ? sanguinicollis *Reiche, Spin.* 125.

Philippine Islands.

C. GRACILIS *White.*

(See Appendix.)

N. India. B. M.
 From Capt. Boyes's Collection.

C. PRINSEPII *White.*

(See Appendix.)

N. India. B. M.

C. LONGIPENNIS *Westwood.*

(See Appendix.)

N. India. Coll. Westwood.
 E. Ind. House.

- C. PECTINICORNIS** N. America ?
Tillus p. *Klug, Berl. Abhand.*
 1842. 270. t. 2. f. 2.
- TENERUS** *Lap.*
Cylistus Klug.
- T. CYANOPTERUS** *Spin.* Manilla.
 I. 165. t. 8. f. 4.
- T. TERMINATUS** *Spin.* C. G. Hope.
 165. t. 12. f. 1. (*variabilis*).
Tillus t. *Dej. Klug.*
- T. PICTUS** *Lap.* Senegal.
Silb. Rev. Zool. IV. 44.
T. dimidiatus (Dej.)
Spin. 166. t. 11. f. 4.
- T. LINEATOCOLLIS** *Spin.* Senegal.
 167. t. 7. f. 4.
- T. PRÆUSTUS** *Lap.* Java. B. M. ?
Silb. Rev. Zool. IV. 43.
Spin. 167. t. 11. f. 2.
 Philippine Islands.
- T. SIGNATICOLLIS** *Lap.* Java.
Silb. Rev. Ent. IV. 44.
Spin. 168. t. 11. f. 3.
 India (Neemuch). B. M.
 var. ?
- T. BIMACULATUS** *Lap.* Senegal.
Silb. Rev. Ent. IV. 44.
Spin. 169. t. 11. f. 1.
- T. BIFASCIATUS** *Spin.* C. G. Hope.
 169. t. 11. f. 5. (*T. variabilis*).
Tillus bif. *Klug.*
T. capensis Gory.
- T. CINGALENSIS** *White.* Ceylon. B. M.
 (See Appendix.)
- T. ABBREVIATUS** *White.* Australia. B. M.
 (See Appendix.)
- SERRIGER** *Spin. (Rev. Zool. Guér. March, 1841.)*
- S. REICHEI** *Spin.* Mexico. B. M.
 171. t. 12. f. 3. G. R. Waterhouse, Esq.

- S. COFFINI, n. s. Mexico.
(See Appendix.)
Opilus C. *Westwood MS.* Coll. Westwood.
- OMADIUS *Lap.*
- O. INDICUS *Lap.* Java. B. M.
Silb. Rev. Ent. IV. 49.
Spin. 175. t. 13. f. 1. Sumatra.
Not. javanus *Dej.* var. Philippine Islands.
Opilo sumatrensis *Dup.*
Clerus prolixus *Klug, Berl.*
Abh. 1842. p. 287.
- O. TRIFASCIATUS *Lap.* Java. B. M.
Silb. Rev. Ent. IV. 49. Dr. Horsfield.
Spin. I. 176. t. 13. f. 3.
Not. spectrum *Dup.*
Clerus modestus *Klug, Berl.*
Abh. 1842. 288. t. 1. f. 2.
- O. KAMELIANUS *White.* Philippine Islands. B. M.
(See Appendix.) From Mr. Cuming's Collection.
- O. NEBULOSUS *Spin.* Java.
II. 133. t. 15. f. 6.
Clerus neb. *Klug, Berl. Abh.*
1842. p. 288.
- O. BIFASCIATUS *Lap.* Ceylon.
Silb. Rev. Ent. IV. 49.
Spin. I. 177. t. 13. f. 2.
- STIGMATIUM *G. R. Gray.*
- S. CICINDELOIDES *G. R. Gray.* Java. B. M.
Griff. A. K. Ins. I. 376. t. 48. Mus. Children, type described
f. 2. by Mr. Gray.
Spin. I. 179. t. 13. f. 4.
Cler. cic. *Klug, Berl. Abh.* 1842. 289.
Var. Cler. rusticus *Dej.*
- S. GILBERTI, n. s. Port Essington. B. M.
From Mr. Gould's Collection.
- S. NATALENSE *Westwood.* Port Natal. B. M.
(See Appendix.)
- S. SPECULARE *White.* Port Essington. B. M.
(See Appendix.)

- S. RUFIVENTRE* *Westwood*. Assam. Coll. *Westwood*.
(See Appendix.)
- S. ELAPHROIDES* *Westwood*. Ceylon. B. M.
(See Appendix.)
- S. SERICELLUM* *White*. Port Natal.

THANASIMUS *Latr.*

- T. MUTILLARIUS* *Latr.* Europe. B. M.
Gen. Cr. et Ins. II. 271. Dr. *Dowler*.
Spin. I. 185. t. 17. f. 4.
Cler. m. Fab. S. El. 279., *Oliv.*
t. 1. f. 12.
Panz. F. I. G. 31. f. 12.
Klug, Berl. Abh. 1842. 290.
Cler. fasciatus Fourcr. Ins. Par.
I. 135.
Attelabus formicaroides Schrank,
Beitr. 6598.

- T. FORMICARIUS* *Latr.* Europe (Brit. Isles). B. M.
Gen. Cr. et Ins. I. 270.
Curtis, B. E. t. 398.
Spin. I. 187. t. 14. f. 2.
Attelabus form. Linn. S. Nat.
II. 620.
Cler. form. Fabr. S. El. I. 280.
Oliv. t. 1. f. 13.
Klug, Berl. Abh. 1842. 290.
Var. *Cler. femoralis Zetterst.*
F. Lapp. I. 122.
Sturm, D. F. t. 221.
Cler. rufipes Hoppe, Ent. Tasch.
1797. p. 136.
Klug, Berl. Abh. 1842. 292.

- T. DUBIUS.*
Clerus dub. (Fabr. S. El. I.
289.?) *Klug, Berl. Abh.*
1842. 292. (*not Spin.*)

- T. RUFICEPS* *Spin.* N. America. B. M.
I. 189. t. 14. f. 3. Nova Scotia. Lieut. *Redman*.
Cler. erythrocephalus Von St. Martin's F., Albany R.
Winth. Hudson's B. G. *Barnston, Esq.*
Trenton Falls, N. York.
Edw. Doubleday, Esq.; Ent. Club of London.

- T. NUBILUS N. America. B. M.
T. abdominalis Kirby, Rev. W. Kirby.
Faun. Bor. Amer.
T. dubius, mas? Barnston. St. Martin's Falls, Albany
Cler. nubilus Klug, *Berl. Mém.* River. G. Barnston, Esq.
1842. 386..
- T. UNDATULUS. U. States (New Hampshire).
Cler. und. Say, *Boston Journ.*
Nat. H. i. 163. (May, 1835.)
- T. MITIS Spin. C. G. Hope.
t. 16. f. 1.
Cler. mi. Klug, *Berl. Abh.* 1842.
T. Verreauxii Spin. i. 90.
- T. QUADRIMACULATUS Spin. Germany (Tyrol). B. M.
i. 192. t. 15. f. 3. M. Chevrolet.
Cler. q. Fabr. *S. El.* i. 281.
Panz. F. I G. 95. 15.
- T. ABDOMINALIS Spin. E. Indies.
t. 15. f. 1.
Cler. abd. Germ. *Col.* 80.
Klug, *Berl. Abh.* 1842. 308.
(nec Chevr.)
T. pictus Spin. i. 194.
- T. SUBSIMILIS, n. s. ——— ? B. M.
- T. INDICUS. C. G. Hope.
Notoxus in. Fabr. *S. El.* i. 288.
- T. COLUMBICUS Spin. Columbia (Bogota).
i. 18. f. 4. ♂ ♀
Cler.? annulipes Buq. ♂
- T. MARMORATUS Spin. C. G. Hope. B. M.
t. 15. f. 2.
Notoxus chinensis Fabr. *S. E.*
i. 288.
T. capensis Spin. i. 197.
Not. marmoratus Klug, *Berl.*
Abh. 1842. 308.
- T. VERSICOLOR Chevr. C. G. Hope.
Rev. Zool. Cuv. 1842. 276.
- T. OBSOLETUS Chevr. C. G. Hope.
Rev. Zool. Cuv. 1842. 276.

- T. APICALIS** *Chevr.* C. G. Hope.
Rev. Zool. Cuv. 1842. 276.
- T. FASCIOLATUS** *Chevr.* C. G. Hope.
Rev. Zool. Cuv. 1842. 276.
- T. ? VIRESCENS** *Chevr.* C. G. Hope.
Rev. Zool. Cuv. 1842. 276.
- T. ? LATUS** *Chevr.* C. G. Hope.
Rev. Zool. Cuv. 1842. 277.
- T. CONFUSUS** *Newman.* S. Australia (Port Philip).
Entom. 364.
- T. ACERBUS** *Newman.* S. Australia (Port Philip).
Entom. 364.
- Fossil.*—**CLERUS ADONIS** *Heer.* Tertiary formation. Ceningen.
Nouv. Mém. Soc. Helv.
Sc. Nat. 1847. 152. t. 5.
 f. 3.

NATALIS *Laporte.*

- N. PORCATA** *Spin.* N. Holland. B. M.
 I. 201. t. 16. f. 2.
Not. porcatus Fabr. S. E. V. D. Land.
 I. 287. Gen. T. Hardwicke.
 Cler. p. *Oliv.* t. 2. f. 17. Earl of Derby.
 Ent. Club of London, &c
- N. CRIBRICOLLIS** *Spin.* N. Guinea.
 I. 203. t. 16. f. 4.
- N. LAPLACEI** *Lap.* "Chili." (?) B. M.
Silb. Rev. Ent. IV. 41. N. Holland.
Spin. I. 204. t. 16. f. 3. Sir John Richardson, M. D.
Not. punctata Gory.

THANEROCLERUS *Lefebvre.* (*Spin.* March, 1841).

- T. BUQUETII** "*Westw.*" E. Indies.
Spin. I. 207. t. 22. f. 2.
 Cler. B. *Lefebvre, Ann. Soc.*
Ent. Fr. v. 577.
 Cler. *sanguinolentus Dej.*
 Var. Cler. *pondicherianus Dup.*

- T. SANGUINEUS *Spin.* N. America. B. M.
 I. 208. t. 17. f. 2. Ent. Club of London.
 Cler. sang. *Say, Bost. Journ.*
Nat. Hist. I. 162.
 Var. Cler. rufipennis *Newm.*
- T. ACCINCTUS. Australia (Port Philip).
 Thanas. accinctus *Newm. Entom.* 364. V. D. Land. Coll. Westwood.
 Var. Cler. viduus *Westw. MS.*
 Cler. medianus *Westw. MS.*
- T. ? DERMESTOIDES. Arabia.
 Cler. derm. *Klug, Berl. Abh.*
 1842. 310.

TROGODENDRON *Guérin.* (March, 1841.)
Xanthoceros Newman (Aug. 1842).

- T. FASCICULATUM *Guér.* N. Holland. B. M.
Icon. R. A. teste expl. III. 53. Sir John Richardson, M. D.
 (1829—1844).
Spin. I. 212. t. 18. f. 1.
 Cler. fasc. *Schreib. Linn. Trans.*
 VI. 195. t. 20. f. 6.
Xanthoceros fas. Newman, Entom. p. 363.
- T. HONESTUM. Australia (Kangaroo I.). B. M.
 Cler. honestus *Newm. Ent.* 16.
 T. fasciculatum var. ?

ERYMANTHUS *Klug.*

- E. GEMMATUS *Klug, Berl. Abhand.* C. G. Hope.
 1842. 327. t. 1. f. 6.
Spin. II. 61. t. 41. f. 5.

NOTOXUS *Fabr.*

- N. TRISTIS *Spin.* Madagascar.
 t. 19. f. 2.
Opilus tris. Klug, Berl. Abh.
 1842. p. 324.
N. funebris Dup. Spin. I. 215.

- N. BUQUETH *Spin.* Africa.
I. 217. t. 16. f. 5.
- N. DORSALIS. Algeria.
Opilus d. Lucas, Expl. Sc. de l'Algérie, t. 20. f. 3.
- N. GIGAS *Lap.* Senegal. B. M.
Silb. Rev. Ent. IV. 42. Var. ?
Spin. I. 218. t. 19. f. 1.
Not. dorsalis *Dej.*
- N. TROPICUS. Sennaar.
Opilus trop. Klug, Berl. Abh.
1842. 322.
Not. gigas *Spin.*
- N. OBSCURUS *Klug.* S. Africa.
Opil. obs. Klug, Berl. Abhan.
1842. 323.
- N. DREGEI *Gory,* C. G. Hope.
Spin. I. 220. t. 19. f. 3.
- N. MOLLIS *Fabr.* Europe. B. M.
S. El. I. 283. Asiatic coasts of Mediter-
Panz. F. I. G. v. 5. ranean.
Spin. I. t. 19. f. 4.
Attelabus m. Linn. Faun. Suec.
186.
Clerus Geoffr. Ins. I. 305.
Opilus m. Latr. Gen. Cr. I. 271.
Samouelle, Ent. Comp. t. 12.
f. 1. var.
Spry & Shuckard, t. 52. f. 3.
Waterhouse, Trans. Ent. Soc.
Lond. I. t. 5. f. 1. (larva).
- N. DOMESTICUS *Sturm.* Germany, France.
Not. subfasciatus *Dej.*
Not. domesticus *Sturm, F. G.*
XI. 16. t. 229. f. n.
Not. unifasciatus *Dahl, Spin.* t.
19. f. 5.
- N. TÆNIATUS. Dalmatia.
Opilus tæniatus Klug, Berl.
Abh. 1842. 320.
Tillus transversalis Parreyss.
Not. flavicornis *Ziegler.*

- N. PALLIDUS.** Italy, France.
Not. centromaculatus Christofori.
Not. pallidus Oliv. t. 1. f. 11.
Spin. t. 8. f. 6.
- N. FRONTALIS** *Spin.* Levant.
t. 27. f. 6.
Opilus fron. Klug, Berl. Abh.
1842. 321.
N. cruentatus Dup. Spin. I. Turkey in Europe.
225.
Not. syriacus Sol.
Cler. thoracicus Friv.
Opilus tho. Klug, Berl. Abh.
1842. 321.
- N. CONGRUUS** S. Australia (Port Philip).
Opilus cong. Newm. Entom. 365. B. M.
Opilus femoralis Westw. MS.
- N. GERMANUS** *Chevr.* N. Germany.
Ann. Soc. Ent. Fr. 1843. p. 37.
- N. SUBFASCIATUS** *Westw. MS.* E. Indies. B. M.
(See Appendix.)
- N. CASTANEIPENNIS** *White.* E. Indies (Bengal).
(See Appendix.) Coll. Saunders.
- N. APICALIS** *White.* Australia. Coll. Parry.
(See Appendix.)
- N. UNICOLOR** *White.* E. Indies.
(See Appendix.) Coll. E. India House
- N. RUFICORNIS** *Westw.*
(See Appendix.) Coll. Westwood.
- N. MÆRENS** *Westw.* S. Australia. Coll. Westwood.
(See Appendix.)
- N. EPHIPPIGER** *White.* Australia. B. M.
(See Appendix.)
- N. PULCHER** *White.* V. D. Land. B. M.
(See Appendix.) Rev. T. Ewing.
- OLESTERUS** *Spinola.* (Rev. Zool. March, 1841.)
- O. AUSTRALIS** *Spinola.* N. Holland (Swan River).
I. 229. t. 20. f. 2.

CLERUS *Fabr.*

- C. LEVIGATUS *Dup.* Mexico. B. M.
Spin. I. 238. t. 21. f. 1.
 Var. Cler. griseo-pilosus *Dup.* Columbia.
 Cler. nebulosus *Buq. Spin.* t. From the Collection of M. Goudot.
 21. f. 2.
 Cler. nigricans *Dup.*
 Cler. brunnipes *Buq.*
 Cler. nitidus *Chevr. Mag. de Zool.* 1843. 24.
- C. FLAVO-SIGNATUS *Dej.* Brazil. B. M.
Spin. I. 239. t. 21. f. 3. Coll. Miers.
- C. BILOBUS *Spin.* Brazil.
 I. 240. t. 21. f. 4.
- C. MINIATUS *Spin.* Brazil. B. M.
 II. 139. t. 47. f. 2. Coll. Miers, Parry.
- C. SOBRINUS *Lap.* Brazil. B. M.
Silb. Rev. Ent. IV. 45.
Spin. I. 241. t. 22. f. 4.
 Cler. rubripes *Dej.*
- C. ERYTHROPUS *Klug, Berl. Abhan.* Brazil.
 1842. p. 304.
- C. COMPTUS *Klug, Berl. Abh.* 1842. Brazil.
 p. 304.
- C. PLANO-NOTATUS *Lap.* Brazil. Coll. Miers, Parry.
Silb. Rev. Zool. IV. 45.
Spin. I. 242.
 Cler. pulchellus *Klug, Berl. Abh.*
 1842. 303., *Spin.* t. 22. f. 2.
 Cler. luctuosus *Dej.*
 C. obliquefasciatus *Chevr. Ann. Soc. Ent. Fr.* 1843. p. 32.
- C. ARTIFEX *Dej.* Cayenne, Para. B. M.
Spin. I. 243. t. 22. f. 3.
- C. DISTINCTUS *Dej.* Brazil. Coll. Parry.
Spin. I. 244. t. 22. f. 6.
 Cler. variegatus *Mannerh.*

- C. PUSILLUS* *Klug, Berl. Abh.* 1842. Brazil. B. M.
 306., *Spin.* t. 22. f. 1.
Cler. columbianus *Dup.*
C. arcuatus *Dej. Spin.* i. p. 245. Columbia.
- C. MYSTICUS* *Klug.* B. M.
Berl. Abh. 1842. 300.
Spin. t. 22. f. 5.
C. antiquus *Dej. Spin.* i. 246. Brazil.
- C. PHALERATUS* *Klug.* Brazil.
Berl. Abh. 1842. 300.
- C. CRABRONARIUS* *Dej.* N. America. B. M.
Spin. i. 247. t. 23. f. 1. Ent. Club of London.
- C. MEXICANUS* *Lap.* Mexico.
Silb. Rev. Ent. iv. 44.
Spin. i. 248. t. 27. f. 2.
Klug, Berl. Abh. 1842. 299.
Cler. assimilis *Bug.*
Cler. Lesueurii *Dup.*
- C. VENATOR* *Chevr. Mag. de Zool.* Mexico.
 1843. p. 18. Guatemala. B. M.
Spin. t. 23. f. 2. (mexicanus *B.*) Edw. Doubleday, Esq.
- C. ANNULATUS* *Eschsch.* Brazil. B. M.
Ent. 50.
Klug, Berl. Abh. 1842. 300.
Cler. myops *Dej.*
C. variegatus *Dej. Spin.* i. 250. Brazil.
 t. 23. f. 4.
- C. DECUSSATUS* *Klug.* Mexico. B. M.
Berl. Abh. 1842. 296.
Spin. t. 25. f. 1
C. Hopfneri *Dej. Spin.* i. 256. From Mr. Argent's Collection.
- C. VERSICOLOR* *Lap.* Brazil. B. M.
Silb. Rev. Zool. iv. 45.
Spin. i. 251. t. 26. f. 6.
Blanchard, Voy. D'Orbigny,
 vii. 89.
Cler. basalis *Dej.* John Miers, Esq.
Cler. nobilis *Dup.*
Var. Cler. cruentatus *Dup. Spin.*
 t. 26. f. 5

- Var. *Cler. scenicus Klug, Berl. Abh.* 1842. 295.
- Var. *Cler. jucundus Klug, Berl. Abh.* 1842. 296.
- Var. *Cler. miniaceus Blanch. l. c. t. 6. f. 1.*
- C. MINUTUS** *Blanch.* S. America, I. los Pajaros,
D'Orbigny, Voy. à l'Amér. Mér. in R. Parana, between B.
 VI. 90. t. 6. f. 4. Ayres and Corrientes.
- C. versicolor** var.
- C. NIGRIVENTRIS** *Blanchard.* S. America, Corrientes.
D'Orbigny, Voy. à l'Amér. Mér.
 VI. 90. t. 6. f. 3.
- C. TRIPLAGIATUS** *Blanchard.* Brazil (Rio).
D'Orbigny, Voy. à l'Amér. Mér.
 VI. 90. t. 6. f. 3. ("trifascia-
 tus" *Blanch. alt. nec Sayi.*)
- C. ICHNEUMONEUS** *Fabr.* N. America. B. M.
Gen. Ins. Mant. 230. Ent. Club of London.
Spin. t. 24. f. 3.
Klug, Berl. Abh. 1842. 294.
- Cler. rufus Oliv. t. 1. f. 16. Spin.* E. Florida. E. Doubleday, Esq.
 I. 252. Georgia. F. Walker, Esq.
- C. BOMBYCINUS** *Chevr.* Mexico.
Col. Mex. 1. fasc.
Guérin, Icon. R. A. t. 15. f. 13.
Spin. t. 24. f. 1.
Klug, Berl. Abh. 1842. 295.
- C. æneicollis** *Déj. Spin.* I. 254.
Cler. purpureus Sturm.
- C. LUNATUS** *Sturm.* N. America. B. M.
Cat. 120. E. Florida.
Klug, Berl. Abh. 1842. 294. E. Doubleday, Esq.
Spin. I. 255. t. 24. f. 2. Ent. Club of London.
- Cler. rufulus Dej.*
- C. ZONATUS** *Klug.* Mexico.
Berl. Abh. 1842. 297.
- Cler. abdominalis Chevr. Col. du Mex.* 1. fasc. (not Germ.)
- C. COLOMBIÆ** *Spin.* Columbia.
 II. 135. t. 46. f. 6.

- C. VARIUS* *Klug.* Mexico.
Berl. Abh. 1842. 296.
- C. OENATUS* *Dup.* Mexico.
Spin. I. 257. t. 25. f. 2.
- C. LAPORTEI* *Guér.* Mexico.
Icon. R. A. texte.
Spin. I. 258. t. 26. f. 4.
Cler. imperialis Dup.
- C. NIGROCINCTUS* *Klug.* Mexico.
Berl. Abh. 1842. 294.
Klug, Berl. Abh. 297.
 Var. *Cl. bicinctus Spin.* I. 260.
 t. 24. f. 4.
- C. RUFIGOLLIS* *Lap.* Cayenne.
Silb. Rev. Zool. II. 45.
Klug, Berl. Abh. 1842. 301.
Spin. I. 260. t. 26. f. 2.
Cler. histrio Dej.
- C. LÆTUS* *Klug.* Mexico.
Berl. Abh. 1842. 301.
- C. TRICOLOR* *Lap.* Mexico.
Silb. Rev. Ent. IV. 46.
Spin. I. 262. t. 24. f. 6.
- C. CINEREO-PILOSUS* *Blanch.* Brazil (Rio Janeiro).
D'Orbigny, Voy. à l'Amér.
Mér. VI. 91.
- C. NIGRIPES* *Say.* N. America. B. M.
Journ. Acad. Nat. Sc. Phil. Ent. Club of London.
 III. 191.
Klug, Berl. Abh. 1842. 293.
Spin. I. 263. t. 25. f. 3.
 Var. *Cler. rufiventris Dej.*
Spin. t. 23. f. 3.
 Var. *Cler. trifasciatus Say.*
Journ. Acad. Nat. Sci. Phil.
 III. 175.
Klug, Berl. Abh. 1842. 293.
- C. ROSMAREUS* *Say.* N. America.
Journ. Acad. Sc. Phil. III. 190.
Klug, Berl. Abh. 1842. 293.

- C. FISCHERI *Spin.* N. Persia.
 I. 265. t. 25. f. 6.
 Cler. dubius *Fisch.* (not Fabr.)
- C. BREVICOLLIS *Kunze.* Hungary.
Spin. I. 266. t. 25. f. 5.
- C. OCULATUS *Dej.* N. America (Mexico). B.M.
Spin. I. 267. t. 26. f. 1.
 From Mr. Argent's Collection.
- C. TROGOSITOIDES *Spin.* Mexico.
 I. 269. t. 27. f. 1.
 Cæstron Trog. *Dup.*
- C. SPHEGEUS *Fabr.* N. America (Rocky Mo.).
S. El. I. 280.
Oliv. t. 1. f. 14.
Spin. I. 270. t. 27. f. 4.
Klug, Berl. Abh. 1842. 298.
- C. ARACHNODES *Klug.* N. America, Mexico.
Berl. Abh. 1842. 299.
Spin. II. 137. t. 3. f. 2.
- C. LUSCUS *Klug.* Mexico.
Berl. Abh. 1842. 299.
- C. VIDUUS *Klug.* N. America, Mexico.
Berl. Abh. 1842. p. 297.
Spin. t. 27. f. 3.
 C. erythrogaster *Sturm. Spin.*
 I. 272.
- C. MÆSTUS *Klug.* Mexico.
Berl. Abh. 1842. 298.
- C. SIGNATUS *Klug.* Mexico.
Berl. Abh. 1842. 301.
Spin. t. 23. f. 5.
 C. bisignatus *Dej. Spin.* I. 272.
 Var. Cler. vulneratus *Klug,*
Berl. Abh. 1842. 302.
- C. LONGULUS *Spin.* Chili.
 II. 138. t. 46. f. 8.
- C. TIBIALIS *Klug.* Brazil.
Berl. Abh. 1842. 302.

- C. **LEPIDUS** *Klug.* Brazil.
Berl. Abh. 1842. 302.
- C. **MACULICOLLIS** *Sturm.*
Spin. I. 273. t. 23. f. 6.
 (Systemoderes?)
- C. **THORACICUS** *Say.* N. America. B. M.
Spin. I. 274. t. 26. f. 3. Ent. Club of London.
 Edw. Doubleday, Esq.
- C. **GAMBIENSIS** *Lap.* Senegal.
Silb. Rev. Ent. IV. 46.
Spin. I. 276. t. 24. f. 5.
- C. *amœnus* *Dej.*
- C. **SILBERMANNI** *Chevr.* Mexico.
Mag. de Zool. 1843. p. 21.
- C. **QUADRINOTATUS** *Chevr.* Mexico.
Mag. de Zool. 1843. p. 22.
- C. **ASSIMILIS** *Chevr.* Mexico.
Mag. de Zool. 1843. p. 17.
- C. **NIGRO-MACULATUS** *Chevr.* Mexico.
Mag. de Zool. 1843. p. 20.
- C. **ZEBRA** *Chevr.* Mexico.
Mag. de Zool. 1843. p. 23.
- C. **SERICANS** *Westw.* Mexico. Coll. Westwood.
 (See Appendix.)
- C. **HOLOSERICEUS** *White.* California. B. M.
 (See Appendix.)
- C. **LATICINCTUS** *White.* Brazil. Coll. Parry, Miers.
 (See Appendix.)
- C. **GUTTULUS** *White.* Australia. B. M.
 (See Appendix.)
- C. **LACORDAIREI** *Spin.* Brazil.
 II. 140. t. 36. f. 6.
- C. **CRASSUS** *Newman.* Australia (Sydney). B. M.
Entom. 15.
- C. **CRUCIATUS** *Macleay.* Australia.
King's Narrative, II. 442.

C. ANTHICOIDES *Westw.* Ceylon. Coll. Westwood.
(See Appendix.)

PYLUS *Newman* (January, 1841).

Yliotus Spinola.

Yliotis Spin. Rev. Zool. Cuv. March, 1841.

P. FATUUS *Newm.* N. Holland (V. D. Land). B. M.
Entom. 364.
Clerus? fat. *Newman, Ent.* p. 35. Ent. Soc. of London.
Yl. Passerinii *Spin.* I. 283.
Yl. ochropus *Spin.* t. 28. f. 3.
Natalis punctata Lap. Gory, Coll.

P. BICINCTUS *Newman.* S. Australia (Port Philip).
Entomologist, 364.

P. ANTHICIDES *Newman.* N. Holland (Port Philip).
Entom. p. 402.

ZENITHICOLA *Spinola.* (*Rev. Zool.* March, 1841.)

Z. AUSTRALIS *Spin.* N. Holland. B. M.
I. 286. t. 28. f. 2.

Z.? OBESA. N. Holland. B. M.
Cler. obesus *Newman.* Ent. Club of London.
White in Stokes's Voyage, app.
. t. 1. f. 9.
Var. obesula.

Z.? FULGENS *Chevr.*
Ann. Soc. Ent. Fr. 1843. p. 41.

PLATYCLERUS *Spinola.* (*Rev. Zool.* March, 1841.)

P. PLANATUS *Spin.* Madagascar.
I. 334. t. 28. f. 4.
Cler. plan. *Lap. Silb. Rev. Ent.*
IV. 4.
Cler. depressus *Dup.*
Cler. madagascariensis *Dej.*

P. ELONGATUS *Spin.* Madagascar.
II. 145. t. 46. f. 2.

TARSOSTENUS *Spinola*.

- T. UNIVITTATUS *Spin.* S. Europe (Tuscany), B. M.
 I. 288. t. 32. f. 3. Brit. Islands.
 Cler. univ. *Rossi, Faun. Etr.*
Mant. I. 44.
 Til. unif. *Curtis, B. E.* t. 267.
 Opi. fasciatus *Steph. Ill. Brit.*
Ent. III. 324.

EBURIPHORA *Spinola* (March, 1841).

- E. CALLOSA *Spin.* Madagascar.
 t. 20. f. 3.
 Opi. callosus *Klug, Berl. Abh.*
 1842. p. 32.
 E. Reichei *Spin.* I. 291.
 E. ? PATRICIA. Madagascar
 Opi. patricius *Klug, Berl. Abh.*
 1842. 325.

TRICHODES *Fabr.*

- T. OCTOPUNCTATUS *Fabr.* S. Europe. B. M.
S. El. I. 208. F. Walker, Esq.
Spin. I. 297. t. 29. f. 2.
 Cler. oct. *Oliv.* t. 1.
 T. UMBELLATARUM *Schönh.* N. Africa (Barbary). B. M.
Syn. II. 49.
Spin. I. 289. t. 29. f. 3.
Klug, Berl. Abh. 1842. 336.
 Cler. umb. *Oliv.* t. 1. f. 2.
 T. DAHLII *Dej.* Italy, Spain.
Spin. I. 300. t. 29. f. 4.
 Trich. affinis *Dahl.* Barbary.
 T. ALVEARIUS *Fabr.* Europe (Brit. Isles). B. M.
S. El. I. 284.
Klug, Berl. Abh. 1842. 334.
Spin. I. 301. t. 29. f. 5.
 Cler. alv. *Panz. F. I. G.* 31. 14. Egypt. Sir J. G. Wilkinson.
Curtis Brit. Ent. I. f. 44.
Steph. Ill. Brit. Ent. III. 326.
Reaum. Mém. VI. t. 8. f. 9—10.
Geoffr. Ins. I. t. 5. f. 4.

- T. AFFINIS** *Dej.* Egypt.
Spin. I. 302. t. 29. f. 6.
- T. APIARIUS** *Fabr.* Europe (Brit. Isles). B. M.
S. El. I. 284. F. Walker, Esq.
Herbst. IV. t. 41. f. 11.
Spin. I. 305. t. 30 f. 2.
Attelabus ap. *L. Syst. Nat.* II. 620.
Cler. ap. *Oliv.* t. 1. f. 4.
Panz. F. I. G. 31. f. 13.
Steph. Ill. Brit. Ent. III. 325.
t. 19. f. 4.
Shuckard & Spry, Brit. Ent.
Col. del. t. 52. f. 5.
Var. *Trich. apicida Ziegl.*
Var. *Trich. arcuatus Bead. Laf.*
Spin. t. 30. f. 2. B.
Var. *Trich. pannonicus Ziegl.*
Trich. interruptus Meg. Spin. t.
30. f. 1.
Trich. subtrifasciatus Sturm.
Var. *Trich. unifasciatus Dahl.*
Trich. corallinus Fald. Mém. Nat.
Mosc. IV. 208.
Var. *Trich. elegans Dej. Spin.* t.
30. f. 2. E.
- T. APIVORUS** *Germ.* N. America. B. M.
Ins. Sp. Nov. I. 81. F. Walker, Esq.
Klug, Berl. Abh. 1842. 332. Ent. Club of London.
Spin. I. 307. t. 30. f. 4.
Trich. trifasciatus Sturm, Cat. t. 1.
n. 6.
Trich. cribripennis Dej.
Trich. Nuttalli Dup.
- T. CRABRONIFORMIS** *Fabr.* Sicily, Corfu. B. M.
S. El. I. 285. Rev. C. Kuper.
Spin. I. 309. t. 30. f. 3. Greece. W. W. Saunders, Esq.
Klug, Berl. Abh. 1842 329. Turkey.
Cler. cr. Oliv. t. 1. f. 1. Asia Minor.
Trich. gulo Parreyss. Syria, Crimea.
Cler. lepidus Brullé, Exp. Sc.
Mor. t. 37. f. 7.
Trich. Zebra Falderm. Mém. Nat.
Mosc. IV. 207. t. 7. f. 3.
Pachyscelis crabron. Hope, Col.
Man. III. 139.

- T. LAMINATUS *Chevr.* Anatolia.
Ann. Soc. Ent. Fr. 1843. p. 40.
- T. ANGUSTUS *Chevr.* Amadan.
Ann. Soc. Ent. France, 1843.
 p. 41.
- T. NOBILIS *Klug.* Turkey (Constantinople). B. M.
Berl. Abh. 1842. 335.
Spin. t. 30. f. 5.
 T. sanguineo-signatus *Dej. Spin.* Greece.
 I. 311.
 Trich. affinis *Dup.* Levant.
 Var. Trich. distinctus *Dej.* W. Persia.
 Trich. illustris *Fald.*
 Trich. variabilis *Dup. Spin.* t.
 30. f. 5. D.
- T. LAFERTEI *Chevr.* Turkey (Constantinople).
Ann. Soc. Ent. Fr. 1843. p. 39.
- T. FAVARIUS *Sturm.* S. Europe (Mediterranean Islands). B. M.
D. F. 11. t. 230.
Klug, Berl. Abh. 1842. 332.
Spin. I. 313. t. 31. f. 1.
Cler. fav. Illig. Mag. I. 80.
Brullé, Expéd. Sc. Mor. t. 37.
 f. 8.
 Trich. senilis *Koll.*
 Var. Trich. punctatus *Stev. Bull.*
Nat. Mosc. I. t. 2. f. 3.
 Trich. insignis *Steven. Bull. Nat.*
Mosc. I. t. 2. f. 2.
 Trich. obliquatus *Brullé, Exp.*
Morée, p. 155. t. 33. f. 9.
 Trich. Phedinus *Godet. Spin.* I.
 t. 31. f. D.
 Trich. hispanicus *Dup.*
 Trich. vicinus *Dej. Spin.* t. 31.
 f. 1. F.
 Trich. axillaris *Dup.*
 Trich. affinis *Dup. Spin.* t. 29.
 f. 1.
 Trich. elegantulus *Dup.*
 Trich. latifasciatus *Spin. Oliv.*

- T. CARCELII *Chevr.* Anatolia.
Ann. Soc. Ent. Fr. 1843. p. 39.
- T. VIRIDI-FASCIATUS *Chevr.* East.
Ann. Soc. Ent. France, 1843.
 p. 40.
- T. SYRIACUS *Dej.* Syria.
Spin. i. 316.
Trich. Olivieri Spin. t. 30. f. 6.
- T. OLIVIERI *Chevr.* Persia.
Guérin, Icon. R. A. t. 16. f. 16.
 p. 154. (non Klug ?)
- T. NUTTALLI *Say.* N. America (near B. M.
Spin. i. 317. t. 31. f. 2. Missouri).
Clerus Nuttalli Kirby, Linn. Trans. Ent. Club. of London.
 xii. 394.
Trich. humeralis Dej.
- T. LEUCOPSIDEUS *Schönh.* Spain. B. M.
Syn. Ins. ii. 49.
Spin. i. 318. t. 31. f. 3. S. France.
Cler. leuc. Oliv. t. 1. f. 6. From Mr. Children's Coll.
Var. Trich. ceraceus Hoffm.
Trich. fuscicornis Spin. (olim).
- T. AULICUS *Dej.* C. G. Hope.
Spin. i. 320. t. 31. f. 4.
Trich. Dejeanii Drege.
- T. AMMIOS *Fabr.* S. Europe. B. M.
S. E. i. 284.
Spin. i. 322.
Cler. amm. Oliv. t. 1. f. 3. N. Africa.
Var. Trich. arthriticus Chevrol. Levant.
Spin. t. 32. f. 1. A.
Trich. dauci Villa Spin. t. 32. Shores of Black Sea.
 f. 1. B.
Trich. flavo-cinctus Dahl. Spin. N. & W. Persia.
 t. 32. f. 1. C.
Trich. smyrnensis Dup. S. & E. Russia.
Trich. vysnagæ Friv. Spin. t. 32. Kirguises Steppes, Siberia.
 f. 1. E.
Trich. omoplatus Dup.
Cler. sipylus Fabr. Ent. Syst. i.
 208.
Oliv. t. 1. f. 7.

Trich. subfasciatus *Fald.*
 Trich. quadriguttatus *Ménétr.*
 Trich. quadripustulatus *Dej. Brullé.*
 Trich. quadripunctatus *Friv.*

Ent. Club of London.
 Prof. E. Forbes.

T. BIFASCIATUS *Fabr.*

S. E. I. 283.

Spin. I. 325. t. 32. f. 2.

Cler. bif. *Oliv.* t. 1. f. 9.

Var. ?

California. B. M.
 From Mr. Hartweg's Collect.

T. DOUGLASIANUS *White.*

(See Appendix.)

California. B. M.

T. HARTWEGIANUS *White.*

(See Appendix.)

California. B. M.

T. ORNATUS *Say.*

Spin. I. 327. t. 31. f. 5.

N. America (U. States).

T. BIZONATUS *Blanchard, D'Orbigny, Voyage à l'Amér. MÉR. VII.*
 p. 89. belongs to the family Dasytidæ, and is related to the
 genus *Danacea Laporte.*

AULICUS *Spinola* (March, 1841).

A. NERO *Spin.* I. 330. t. 27. f. 5. Mexico.

A. INSTABILIS *Spin.* N. Holland. B. M.

I. 331. t. 28. f. 1.

Cler. inst. *Newm. Ent.* p. 15.

Cler. auratus *Gory.*

Gen. Th. Hardwicke.

Var. Aul. episcopalis *Spin.* (olim).

Ent. Club of London.

Var. Cler. tibialis *Westw. MS.*

Var. Cler. castanipes *Westw. MS.*

A. PANTOMELAS.

Notoxus pan. *Boisd. Voy. de N. Zealand.* B. M.

l'Astrol. t. 6. f. 14.

White, Voy. Erebus and Terror,

W. Colenso, Esq.

Ins. N. Zeal. p.

Var. C. longulus *Spin.* ? t. 46. f. 8.

MUISCA *Spinola.*

M. BITÆNIATA *Spin.*

Columbia.

II. 148. t. 47. f. 4.

PHLOIOCOPUS *Guérin. Spin.* 1841., expl. III. 54. (1829. 1844.)

- P. TRICOLOR *Guérin.* Senegal. B. M.
Icon. R. A. texte.
Spin. I. 338. t. 17. f. 1. Collection of Dr. Leach.
 Cler. Lelieurii (*Dej.*) *Dup.*
 Cler. Lesueurii *Dej.*
 Opilus interruptus *Klug, Berl.*
Abh. 1842. 323.

- P. BASALIS Arabia (Mt. Sinai).
 Opilus basalis *Klug, Berl. Abh.*
 1842. 323.

- P. GUERINII *White, n. s.* Africa. B. M.
 an ♀ tricolor.

- P. BUQUETII *Spin.* Senegal.
 I. 340. t. 18. f. 3.
 Not. bipartitus *Bug.*
 Not. dimidiatus *Bug.* (olim).

- P. ? SUBEROSUS. Madagascar.
 Opilus suberosus *Klug, Berl. Abh.*
 1842. 324.

ENOPLIUM *Latr.*

- E. DENTATUM. S. Europe (Italy, B. M.
 Dalmatia).
 Enop. serraticorne *Latr. Hist.*
Nat. Crust. et Ins. 9. 146. t.
 76. f. 9.
Panz. F. I. G. 26. 13.
Spin. I. 345. t. 33. f. 4.
Klug, Berl. Abh. 1842. 359.
 Tillus serraticornis *Hellwig, Rossi,*
Mant. Ins. I. 37.
Oliv. II. t. 1. f. 1.
Fabr. S. E. I. 282.
 Dermestes dentatus *Rossi, Faun.*
Etr. I. 34. t. 3. f. 2.

- E. QUADRIPUNCTATUM *Say.* N. America.
Journ. Acad. Sc. Phil. III. 188.
Amer. Ent. III. t. 41.
Spin. I. 346. t. 34. f. 5.
Klug, Berl. Abh. 1842. 367.

PELONIUM *Spin.*

- P. LAMPYROIDES** *Spin.* Columbia.
 I. 349. t. 33. f. 6.
 Enopl. Lamp. *Dup.*
- P. LUCTUOSUM** *Spin.* Cayenne.
 I. 351. t. 28. f. 6.
 Enopl. l. *Dej.*
- P. SUTURALE** *Spin.* Brazil.
 I. 352. t. 35. f. 2.
 Enopl. sut. *Buq.*
- P. FLAVO-LIMBATUM** *Spin.* Brazil. Coll. Miers.
 I. 354. t. 35. f. 5.
 Enopl. f. *Dup.*
- P. PILOSUM** *Spin.* N. America. B. M.
 I. 356. t. 33. f. 2.
 Lampyris pil. *Forster, Nov. Sp.* Ent. Club of London.
Ins. 49.
 Var. Enopl. cinctum *Dej. Spin.*
 t. 36. f. 4.
 Enopl. marginatum *Say, Journ. Acad. Sc. Phil.* III. 187.
 Enopl. pilosum et E. marginatum
Klug, Berl. Abh. 1842. p. 360.
 Enopl. onustum *Say, Amer. Ent.*
 III. t. 41.
- P. AMÆNUM** *Guér.* South America (interior of
Icon. R. A. texte expl. III. 55. Brazil, Minas Geraes).
 (1829. 1844.)
Spin. I. 358.
 Pelon. fasciculatum *Spin.* Magellania.
 t. 36. f. 3.
 Enopl. amæn. *Guér. Icon. R. A.*
 Enopl. metalloideum *Dup.*
 Enopl. fasciculatum *Klug.*
- P. CYANEO-MACULATUM.** S. America.
 Enopl. c. *Blanchard, D'Orbigny*
Voy. à l'Am. MÉR. VI. 94. t. 6.
 f. 9.
 E. fasciculatum *Klug, Berl. Abh.*
 1842. 364. t. 1. f. 12. ?
 Var. P. amæn. ?

- P. SCOPARIUM.** Brazil.
 Enopl. scoparium *Klug, Berl. Abh.* 1842. 365.
 Var. *P. amœn.* ?
 Enopl. fugax *Klug*, l. c. 365.
 Var. Enopl. leucophæum *Klug*, l. c. 366.
- P. NIVEUM** *Chevr.* S. America (Rio Janeiro).
Ann. Soc. Ent. Fr. 1843. p. 35.
- P. DECORUM.** Brazil.
 Enopl. decorum *Klug, Berl. Abh.* 1842. t. 1. f. 11.
- P. VETUSTUM** *Spin.* N. America (U. States).
 l. 360. t. 35. f. 4.
 Enopl. vetustum *Dej.*
- P. AMABILE** *Spin.* Columbia.
 l. 362. t. 33. f. 1.
 Enopl. am. *Dup.*
- P. MARGINIPENNE** *Spin.* N. America (U. S.). B. M.
 l. 363. t. 35. f. 6.
 Enopl. m. *Dej.*
 Enopl. circumcinctum *Dup.* Ent. Club of London.
 E. Doubleday, Esq.
- P. CRINITUM** *Spin.* W. Indies (St. John's).
 t. 33. f. 7.
 Enopl. crinitum *Klug, Berl. Abh.* S. America (Carthagena).
 1842. 369.
 Pelon. collare *Spin.* l. 364.
 Enopl. coll. *Dej.*
 Enopl. punctato-striatum *Dup.*
 Var. Enopl. flavum *Dup.*
- P. QUADRISIGNATUM** *Spin.* Columbia.
 l. 365. t. 33. f. 3.
 Enopl. q. *Dup.*
- P. HUMERALE** *Spin.* Brazil.
 l. 366. t. 35. f. 3.
 Enopl. h. *Buq.*
- P. SCUTELLATUM** *Buq.* Brazil.
Spin. II. 156.
 Enoplium crinitum *Dej.*

- P. TESTACEUM** *Bug.* Peru. B. M.
Spin. II. 158. t. 47. f. 6.
Enopl. pilosum Klug. Brazil.
- P. VARIABILE** *Spin.* Brazil.
 I. 367. t. 36. f. 5.
Enopl. badium Dup. Columbia.
Enopl. nigricorne Dup.
- P. PRÆEUSTUM** *Spin.* Magellania.
Enopl. præust. Reiche.
Enopl. principale Dup.
- P. SEMINIGRUM** *Chevr.* Columbia.
Ann. Soc. Ent. Fr. 1843. p. 34.
- P. VIRIDIPENNE** *Spin.* Brazil.
 I. 370. t. 35. f. 1.
Enopl. viridip. Kirby, Linn. Tran. Coll. Ent. Soc. London,
 XII. 393. olim Kirby.
Guérin. Icon. R. A. t. 16. f. 20.
Klug, Berl. Abh. 1842. 362.
- P. AURIPENNE.** Brazil.
Opilus auripennis Hope, Trans.
Zool. Soc. Lond. I. 95. t. 13. f. 5.
Enopl. aurip. Klug, Berl. Abh.
 1842. 395.
- P. KIRBYI.** Brazil. B. M.
Lasiodera Kirbyi Gray, Griff. Gen. T. Hardwicke.
A. K. Ins. I. 376. t. 48. f. 3.
Enopl. Kirbyi Klug, Berl. Abh.
 1842. 362.
- P. SEX-NOTATUM.** Brazil. B. M.
Enopl. sex-notatum Klug, Berl.
Abh. 1842. 368.
- P. NIGROSIGNATUM** *Spin.* Columbia. B. M.
 I. 371. t. 34. f. 6. From Goudot's Collect.
Enopl. nigros. Dup.
Enopl. variegatum Spin. (olim.)
- P. BUQUETHI** *Spin.* Columbia.
 125. t. 46. f. 3.
- P. GALLERUCOIDES** *Spin.* Cayenne.
 I. 372. t. 39. f. 2.
Enopl. Galler. Dej.

- P. TRIFASCIATUM** *Spin.* Brazil. B. M.
 t. 34. f. 3.
Clerus trifasciatus Laporte, Silb.
Rev. Ent. iv. 47.
Enopl. trifasciatum Klug, Berl.
Abh. 1842. 362.
Pelon. Cleroides Spin. i. 374.
Clerus Spinolæ Germ.
Var. Enopl. rufipes Klug, Berl.
Abh. 1842. 362.
- P. LITURATUM** *Spin.* Brazil.
 i. 376. t. 34. f. 1.
Enopl. lit. Kirby, Linn. Trans.
 xii. 393. Coll. Ent. Soc. Lond.
 olim Kirby.
Enopl. liberatum Lap. Hist. An.
Art. i. 289.
- P. PULCHELLUM** *Spin.* Brazil.
 i. 377. t. 34. f. 4.
Enopl. p. Dej.
Philyra helopioides Lap. Silb.
Rev. Ent. iv. 54.
- P. HIRTULUM** *Spin.* Brazil.
 i. t. 33. f. 5.
Enoplium hirtulum Klug, Berl.
Abh. 1842. 367.
Pelon. cribripenne Spin. i. 379.
- P. APICALE** *Buquet.* Brazil.
Spin. ii. 155. t. 47. f. 4.
- P. VITTATUM** *Buq.* Brazil.
Spin. ii. 159. t. 47. f. 3.
- P. ORNATUM.**
Enopl. ornatum Klug, Berl. Brazil.
Abh. 1842. 363. t. 11. f. 11.
- P. PUNCTATISSIMUM** *Chevr.* N. America.
Ann. Soc. Ent. Fr. 1843. p. 34.
Tillus bicolor Say, Journ. Acad.
Sc. Phil. v. 174.?
- P. FIMBRIOLATUM** *Chevr.* Brazil.
Ann. Soc. Ent. Fr. 1843. p. 35.
- P. TERMINATUM** *Blanch.* Brazil (Rio Janeiro).
Voy. à l'Amér. MÉR. vi. p. 94.

- P. OBSOLETUM *Blanch.* Bolivia (Chiquitos).
Voy. à l'Amér. Mér. vi. p. 95.

APOLOPHA *Spinola.* (*Rev. Zool.* March, 1841.)

- A. REICHEI *Spin.* Columbia.
 i. 383. t. 36. f. 1.

MONOPHYLLA *Spinola.* (March, 1841.)

- M. MEGATOMA *Spin.* N. America. B. M.
 i. 385. t. 28. f. 5.
Enopl. meg. Dej.

- M. TERMINATA *Spin.* N. America.
 ii. 126. t. 6. f. 3.
Tillus (Macrotelus) terminatus
Klug, Berl. Abh. 1842. t. 2. f. 4.
Say, Bost. Journ. i. 160.

PHYLLOBÆNUS *Spinola.*

- P. TRANSVERSALIS *Spin.* N. America.
 ii. 4. t. 40. f. 6.

EPIPHLÆUS *Dej.* (*Spin.* March, 1841.)

- E. 12-MACULATUS. Cayenne.
Enoplium duodecimmaculatum
Klug, Berl. Abh. 1842. 370. Coll. Miers
 E. 12-punctatus *Spin.* t. 42. f. 1.
 E. pantherinus *Dej. Spin.* ii. 8.
- E. PANTHERINUS *Chevr.* (Enopl.) Cayenne.
Ann. Soc. Ent. Fr. 1843. 35.
 Var. E. 12-mac. ?
- E. NUBILUS. Brazil.
Enopl. nubilum Klug, Berl. Abh.
 1842. 370. t. 2. f. 15.
- E. MUCOREUS. Brazil.
Enopl. mucoreum Klug, Berl.
Abh. 1842. 371.
- E. FASCIATUS. Brazil (Para).
Enopl. fasciatum Klug, Berl. Abh.
 1842. 371.
 E. mucoreus ♀ ? *Spin.*

- E. BUQUETHI *Spin.* Cayenne.
II. 10.
- E. ORNATUS *Dup.* Cayenne.
Spin. II. 12.
- E. VARIEGATUS *Spin.* Brazil (Para).
t. 38. f. 3.
Enopl. variegatum *Klug, Berl.*
Abh. 1842. 372.
E. tomentosus *Spin.* II. 13. Cayenne.
- E. SPECULUM. Brazil.
Enopl. speculum *Klug, Berlin.*
Abh. 1842. 372.
- E. BALTEATUS *Chevr.* (Enopl.) Brazil.
Ann. Soc. Ent. Fr. 1843. p. 36.
- E. MARGINELLUS *Spin.* Cayenne.
II. 15. t. 42. f. 2.
Enopl. m. *Buq.*
- E. HUMERALIS *Spin.* Cayenne.
II. 16. t. 38. f. 5.
Enopl. hum. *Buq.*
- E. SERICEUS. Brazil.
Enopl. sericeum *Klug, Berl. Abh.*
1842. 373. t. 2. f. 16.
- E. DISTROPHUS. N. America.
Enopl. distrophum *Klug, Berl.*
Abh. 1842.

PLOCAMOCERA *Spinola.*

- P. SERICELLA *Spin.* Carthagena.
II. 19. t. 38. f. 4.
Epiphyl. ser. *Dup.*

ICHNEA *Laporte.*

- I. LYCOIDES *Laporte.* Brazil. B. M.
Silb. Rev. Ent. iv. 55.
Spinola, II. 22. t. 37. f. 3. Columbia.
Enopl. Thomasii *Dej.* From Goudot's Collection.
Enopl. melanurum *Klug, Berl.*
Abh. 1842. 376. t. 2. f. 12.

Var. *Enopl. æquinoctiale* *Buq.*
Spin. t. 37. f. 4, 5.

Var. *Pyticera lycoides* *Dup. Spin.*
 t. 37. f. 6.

♀ *Enopl. laterale* *Klug, Berl. Abh.*
 1842. t. 2. f. 13.

Var. *Enopl. præustum* *Klug.* l. c.
 t. 1. f. 14.

I. *ENOPLIOIDES* *Spin.*

Cayenne, Surinam,
 Columbia.

II. 25. 163. t. 37. f. 1.
Enopl. opacum *Klug, Berl. Abh.*
 1842. 377.

Enopl. roseicolle *Dej.* ♀

Enopl. crinipenne *Dup.* ♀

Enopl. nigripenne *Buq.* ♀

Enopl. lymexylonoides *Dup.* ♂

Var. ♂ *Enopl. pubescens* *Dej.*
Spin. t. 37. f. 2.

Var. *Enopl. marginellum* *Klug,*
Berl. Abh. 1842. 376.

I. *SUTURALIS.*

Brazil (Bahia).

Enopl. suturale *Klug, Berl. Abh.*
 1842. 378. t. 1. f. 15.

Ichnea enoplioides *Spin.* Var.

I. *ATERRIMA.*

Mexico.

Enopl. aterrimum *Klug, Berl.*
Abh. 1842. 378.

Ichn. Enopl. var. ? Spin.

I. *DIMIDIATIPENNIS* *Spin.*

Columbia.

II. 162. t. 46. f. 1.

I. *DIVISA* *Chevr.*

Brazil.

Ann. Soc. Ent. Fr. 1843. p. 37.

I. *CALCEATA* *Chevr.*

Brazil.

Ann. Soc. Ent. Fr. 1843. p. 37.

EVENUS *Laporte.*

E. *FILIFORMIS* *Laporte, Silb. Rev. Ent.*

IV. 42.

Spin. t. 38. f. 2.

ALLELIDEA *Waterh.*A. CTENOSTOMOÏDES *Waterh.*

N. Holland (Sidney). B. M.

Trans. Ent. Soc. Lond. II. 192.

Charles Darwin, Esq.

PEZOPORUS *Klug (nec Illig.)*.

P. COARCTATUS.

S. Africa (Caffraria).

Clerus coarctatus *Klug, Berl.**Abh.* 1842. 311. t. 1. f. 3.ELLIPOTOMA *Spin.*E. TENUIFORMIS *Spin.*

Columbia.

II. 38. t. 40. f. 5.

Not ten. *Dup.*LEMIDIA *Spinola.*L. NITENS *Spin.*

N. Holland.

B. M.

II. 34. t. 38. f. 1.

V. D. Land.

Hydn. n. *Newm. Entom.* p. 36.*Hombr. and Jacq. Voy. au Pole**Sud*, t. 4. f. 15.Clerus nitens *Klug, Berl. Abh.*

1842. 311. t. 2. f. 8.

Cler. clytoides *Gory.*Cler. scalaris *Dup.*L. INANIS *Germ. Linn.*

Australia (Adelaide).

Ent. III. 137.

HOPLOCLERUS, n. g.

H. BIACULEATUS *Westw. MS.*

S. Australia.

Coll. Westwood.

HYDNOCERA *Newman. (Mag. of Nat. Hist. April, 1838.)*H. BICARINATA *Spin.*

II. 43. t. 39. f. 1.

Cler. curvipennis *Dup. Coll.* ♂Cler. megascelisoides *Dup. Coll.* ♀

H. ATTENUATA.

S. Brazil.

Clerus attenuatus *Klug, Berl.**Abh.* 1842. 312. t. 1. f. 4.

H. LIVIDA.

Brazil.

Clerus lividus *Klug, Berl. Abh.*

1842. 313.

- H. HUMERALIS** *Newm.* N. America, B. M.
Mag. Nat. Hist. n. s. iv. 362. N. Carolina, E. Florida,
Spin. II. 44. t. 39. f. 2. Georgia.
Cler. hum. *Say, Phil. Journ. Nat.* Ent. Club of London.
&c. III. 192.
Germ. Spec. Ins. Nov. t. 80.
Phyll. axillaris Dej. ♂
Var. Phyll. cyaneus Dej.
- H. BASALIS.** Aragua.
Clerus basalis Klug, Berl. Abh.
 1842.
- H. RUFIPES** *Newman.* N. Amer. (U. St.) B. M.
Mag. Nat. Hist. n. s. iv. p. E. Florida.
 363. Ent. Club of London.
 E. Doubleday, Esq.
- H. SERRATA** *Newm.* N. America (Ohio, B. M.
Ent. Mag. v. 380. fig. p. 379. Mt. Pleasant).
Mag. Nat. Hist. n. s. iv. 363. Carthagera.
Spin. II. 47. t. 39. f. 4.
Phyllob. quadrim. Dej. Ent. Club of London.
- H. CINCTA** *Spin.* Cayenne.
 II. 48. t. 39. f. 5.
- H. SUTURALIS** *Spin.* Columbia.
 t. 39. f. 6.
Clerus suturalis Klug, Berl. Abh.
 1842.
H. limbata Spin. II. 49. N. America.
Phyll. limb. Dej. ♂
Phyll. transversalis Dej. ♀
Var. Phyll. æquinoctialis Dej.
- H. TENELLA.** Mexico.
Clerus tenellus Klug, Berl. Abh.
 1842.
- H. CRUCIATA.** Columbia.
Theano cr. Chevr. Ann. Soc.
Ent. Fr. 1843. p. 33.
- H. BRACHYPTERA** *Spin.* N. America.
 t. 40. f. 1.
Clerus brachypterus Klug, Berl.
Abh. 1842. p. 313.
H. lineatocollis Spin. II. 51.
Phyll. l. Dej.

- H. CURTIPENNIS* *Newman*. N. Amer. (U. St.) B. M.
Mag. Nat. Hist. n. s. iv. 364. E. Florida.
 E. Doubleday, Esq.
- H. ÆGRA* *Newman*. N. America. E. Florida.
Mag. Nat. Hist. n. s. iv. 364. E. Doubleday, Esq.
- H. BREVIPENNIS* *Spin.* Columbia.
 II. 46. t. 39. f. 3.
 Cler. brevip. *Dup.*
- H. STENIFORMIS* *Spin.* N. America.
 t. 40. f. 2.
 Clerus stenif. *Klug, Berl. Abh.*
 1842.
H. subænea *Spin.* II. 51.
 Phyll. subæneus *Dej.*
- H. AZUREA* *Spin.* Columbia.
 II. 52. t. 40. f. 3.
 Cler. az. *Dup. coll.*
- H. PUNCTATA* *Spin.* N. America.
 II. 54. t. 40. f. 4.
 Phyll. punct. *Dej.*
- H. ? CONFERTA* *Newman*. S. Australia (Port Philip).
Entom. 365.
- H. ? MALTHINUS* *Newman*. V. D. Land. B. M.
Entom. 37. Ent. Club of London.
- H. PECTORALIS* *Westw. MS.* S. Australia.
 (See Appendix.)
- H. FLAVO-VARIA* *Westw. MS.* Australia. B. M.
 (See Appendix.) (var.)
- H. FLAVO-LINEATA* *Westw. MS.* Australia. B. M.
 (See Appendix.)
- H. TASMANICA*, n. s. V. D. Land.
 (See Appendix.) Coll. Saunders.
- H. PICTA*, n. s. Port Philip.
 (See Appendix.) Coll. Saunders.
- EURYMETOPUM* *Blanchard*.
- E. MACULATUM* *Blanch.* Chili (Valparaiso).
D'Orbigny, Voy. à l'Am. MÉR.
 VI. 92. t. 6. f. 8.

- E. PALLENS *Blanch.* Chili (Valparaiso). B. M.
D'Orbigny, Voy. à l'Am. MÉR.
 VI, 93.

E. macul. var.?

- E. FULVIPES *Blanch.* Chili.
D'Orbigny, Voy. à l'Am. MÉR.
 VI, 93. t. 6. f. 7.

PLATYNOPTERA *Lap.*

- P. DUPONTI *Spin.* Mexico.
 II, 64. t. 41. f. 4.
Enopl. dilatatum Dup. Coll.

- P. GORYI *Lap.* Cayenne.
Silb. Rev. Ent. IV, 54.
Spin. II, 65. t. 41. f. 1.
Enopl. fasciatum Buq.

- P. LYCIDES *Chevr.* Carthagena. B. M.
Silb. Rev. Ent. II, 18.
Spin. II, 67. t. 41. f. 2.
Enopl. lyc. Dej. M. Chevrolat.

- P. VITTICEPS *Blanch.* Brazil (Rio Janeiro).
Voy. à l'Amér. MÉR. VI, 95.

PYTICERA *Dupont.*

- P. DUPONTI *Spin.* Brazil. Coll. Miers.
 II, 71. t. 41. f. 3.
Enopl. flaveolatum Dup.

RYPARUS *Spinola.*

- R. TOMENTOSUS *Spin.* Senegal. B. M.
 II, 76. t. 41. f. 6.
Cler. tom. Dej. J. D. Tebbs, Esq.

PROSYMNUS *Lap.*

- P. CRIBRIPENNIS *Lap.* Senegal.
Silb. Rev. Ent. IV, 51.

LEBASIELLA *Spinola.*

- L. ERYTHRODERA *Spin.* Columbia.
 II, 79.
Leb. lepida Spin. t. 43. f. 1.
Coryn. erythroderus Dup.

- L. MARGINELLA. California?
 C. marg. *Chevr. Ann. Soc. Ent.*
Fr. 1843. p. 42.

ORTHOPLEURA *Spinola.*

- O. DAMICORNIS *Spin.* N. America. B. M.
 II. 80. t. 42. f. 4.
 Tillus damic. *Fabr. S. El.* I. 282.
 Enopl. thoracicum *Say.*
 Var. Enopl. nigripenne *Dup.*
- O. SANGUINICOLLIS *Spin.* Central Europe.
 II. 82. t. 42 f. 5.
 Corynetes s. *Fabr. S. E.* I. 187.
 Enoplum Weberi *Latr. Gen. Cr.*
et Ins. I. 271.
 E. s. *Sturm, Deutsch. F.* XI. t.
 233.
 Tillus Weberi *Fabr. S. E.* I. 282.
- O. QUADRATICOLLIS *Spinola.*
 II. 167. note t. 32. f. 4.

CHARIESSA *Perty.*

- C. RAMICORNIS *Perty.* Brazil. Coll. Miers.
Del. Ins. t. 22. f. 3. Coll. Parry.
Spin. II. 86. t. 45. f. 1.
Klug, Berl. Abh. 1842. 363.
- C. VESTITA *Spin.* Mexico. B. M.
 II. 88. t. 45. f. 2.
 Brachymorphus vestitus *Chevr.*
Col. Mex. 7. 150.
 Enoplum vest. *Klug, Berl. Abh.*
 1842. 363. t. 1. f. 10.
 Corynetes spectabilis *Lap. Silb.*
Rev. Ent. IV. 50.

NOTOSTENUS *Dej.*

- N. VIRIDIS *Spin.* C. G. Hope. B. M.
 II. 91. t. 42. f. 3.
 Anobium viride *Thunb. Nov. Ins.* Gen. Th. Hardwicke.
 I. 9. Dr. A. Smith.

- N. CÆRULEUS. C. G. Hope.
Anobium cæruleum Thunb. Nov.
Ins. I. 10.
Corynetes Thunbergii Klug, Berl.
Abh. 1842. 354.

CORYNETES Payk.

- C. SCABRIPENNIS *Spin.* Senegal.
 II. 94. t. 43. f. 2.
- C. ANALIS *Klug.* Caffraria.
Berl. Abh. 1842. 348.
Spin. t. 43. f. 3.
Cor. pallicornis Spin. II. 95. Senegal.
Cor. abdominalis Fabr. S. E. I.
 186. ?
- C. PECTORALIS *Klug.* Caffraria.
Berl. Abh. 1842. 348. t. 1. f. 7.
- C. VIOLACEUS *Payk.* N. Europe (Brit. Isl.). B. M.
F. Suec. II. 275.
Curtis, Brit. Ent. VIII. t. 351.
Spin. II. 96. t. 43. f. 4. ?
Coryn. chalybæus Dej. ?
- C. RUFICORNIS *Sturm.* S. Germany.
D. F. 42. t. 232. f. p. p.
Klug, Berl. Abh. 1842. p. 347.
C. violaceus var. Spin. ?
- C. PUSILLUS *Klug.* Sardinia.
Berl. Abh. 1842. p. 347.
C. violaceus var. Spin. ?
- C. GENICULATUS *Klug.* Portugal, Spain.
Berl. Abh. 1842. p. 347.
C. violaceus var. Spin. ?
- C. SEMISTRIATUS *Spin.* C. G. Hope.
 II. 98. t. 43. f. 5.
- C. DISCOLOR *Klug.* Mexico (Xalappa).
Berl. Abh. 1842. 353.
- C. PALLIPES *Klug.* Mexico (Xalappa).
Berl. Abh. 1842. 353.

NECROBIA Latr.

- N. RUFIPES *Oliv.* Europe (Brit. Islands). B. M.
 t. 1. f. 2. Asia, Africa, America,
Curtis, Brit. Ent. VIII. (under Australia.
 fol. 350.)
Spin. II. 101. t. 42. f. 6.
Coryn. rufipes Fabr. S. El. I.
 186.
 Var. N. amethystina *Steph. Ill.*
Brit. Ent. V. 417.
- N. RUFICOLLIS *Latr.* Europe (Brit. Islands). B. M.
Gen. Cr. & Ins. I. 274. Asia, Africa, America,
Oliv. t. 1. f. 3. Australia.
Spin. II. 103. t. 43. f. 6.
Curtis, Brit. Ent. VII. fol. 350.
Coryn. rufic. Fabr. S. El. I. 286.
- N. VIOLACEA *Latr.* Europe, Asia, America. B. M.
Gen. Cr. et Ins. I. 274.
Oliv. t. 1. f. 1.
Spin. II. 105. t. 44. f. 1.
Coryn. angustatus Falderm. MS.
Necr. quadra Marsh. 323. 4.
Curtis, VIII. (under fol. 350.)
- N. TIBIALIS *Spin.* C. G. Hope.
 II. 107. t. 44. f. 2.
- N. DEFUNCTORUM *Spin.* Spain (Andalusia).
 II. 108. t. 44. f. 3.
Coryn. def. Walth. Reise. Tyrol,
 II. 63.
Silb. Rev. Ent. IV. 143.
Coryn. carbonarius Dej.
- N. ATRA. S. Africa.
Corynetes ater Klug, Berl. Abh.
 1842. 353.
- N. BICOLOR *Spin.* Spain.
 II. 109. t. 44. f. 4.
Cor. b. Lap. Silb. Rev. Ent. IV.
 52.
Cor. thoracicus Dej.
- N. PINGUIS *Westw MS.* Australia. B. M.
 (See Appendix.)

APPENDIX.

DESCRIPTIONS OF NEW SPECIES.

CYLIDRUS NIGRINUS, n. s.

Epistome somewhat 3-lobed, the middle lobe with three equal crenations. Antennæ with the five basal joints of a clear amber-yellow; the other joints black. Head, thorax, elytra, and greater portion of under parts, of a deep black; at least three of the terminal joints of abdomen, both above and below, of a reddish yellow. First pair of legs black, second and third pairs pale yellow, the base of the tibia in the latter alone being brownish. Head and thorax punctured, and furnished with short hairs; elytra smooth.

Length $3\frac{1}{2}$ lines.

Hab. Van Diemen's Land.

TILLUS HILARIS Westw. n. s.

Head rufous, punctured. Antennæ and palpi fulvous; prothorax chestnut-red, in front blackish, shining, irregularly punctured; elytra at the base, for a third part, of a testaceous rufous, the remainder black with a white band in the middle. Elytra punctato-striate the punctures extending through the white band; the tip of elytra smooth. Body beneath and legs rufous, abdomen black.

Length 2 lines.

Hab. Van Diemen's Land.

TILLUS PICIPENNIS Westw.,

Small, narrow, and of a pitchy brown colour. Head and prothorax punctured, the latter with various smooth spaces. Antennæ of a pitchy chestnut-brown, the three last joints black; elytra pitchy brown, punctato-striate; a wide white band behind the middle; the tips, for a considerable space, smooth. Legs pitchy brown.

Length $1\frac{3}{4}$ line.

Hab. East Indies.

TILLUS BIFASCIELLUS, n. s.

Serrated part of antennæ with eight joints. Head, thorax, abdomen, and elytra of a deep blackish blue. Head and thorax polished, covered, especially on the sides, with longish ashy hairs. Elytra punctato-striate for rather more than half their length, with two transverse yellow bands, the first not far from the base, and not reaching either margin of elytron; the second just beyond the middle, reaching the outer edge of elytron, but not extending to suture, somewhat angulated in the middle. The tip of each elytron closely covered with ashy silk-like hairs; sides of terminal segments of abdomen beneath with close silky hairs.

Length $3\frac{1}{2}$ lines.

Hab. Philippine Islands.

TILLUS DISTINCTUS, n. s.

Thickened part of antennæ very slightly serrated and compressed; reddish brown, the edges black. Head and fore-part of thorax above black; rest of thorax above and below red; base of elytra and base of abdomen beneath red; rest of elytra black, excepting a transverse yellow band beyond the middle; tip of elytra covered closely with yellowish silky hairs. Elytra, as far as commencement of yellow band, rather coarsely punctato-striate. Legs red, femora of hind legs black. Greater part of abdomen beneath black; the pointed part of segment between hind legs red.

Length $2\frac{3}{4}$ lines.

Hab. Java. Mus. East India House.

PRIOCERA MIERSIANA, n. s.

Elytra entire, and conjointly rounded at the end; the upper surface for half their length with longitudinal punctured striæ; those on external margin abbreviated. Antennæ red. Head and thorax of a deep rusty brown: thorax above in the middle somewhat depressed, a transverse grooved line not far from the front, and nearly parallel with it; on each side of the depression in front a roundish smooth tubercle; sides of thorax with several delicate streaks. Base of elytra and suture as far as middle of a brownish red; an irregular branched yellow spot on the base of elytra, an angled yellow spot on lateral edge of elytra before the middle, and a transverse yellow spot beyond the middle near the suture, but not reaching to it; a large deep brown band beyond the middle, and extending up the sides of elytra to

beyond the angled yellow spot; tip of elytra of a pale yellow. Under side of abdomen pitch brown; terminal segment paler. The whole insect is covered with longish pale ochraceous hairs.

Length $4\frac{1}{2}$ lines.

Hab. Brazil. *Coll.* Miers.

XYLOBIUS ALBO-VARIUS Westw.

Elongated, slender, of a dark steel-blue colour, densely punctured. Head between the eyes of a silky white. Prothorax with a frontal band, and two spots on the sides behind of a silky white. Elytra with two lunules at the base and seven spots, the third larger than the rest and common to both elytra, of a silky white; tips truncated. Antennæ very short, and with the legs fulvous.

Length 5 lines.

Hab. E. Indies (Himalaya).

SYSTEMODERES DASYTOIDES Westw. n. s.

Elytra of a dark metallic green with yellowish reflexions, thickly punctured, thickened at the suture. Head and front of thorax black, rest of thorax red; in some specimens the whole of the thorax is black. Antennæ, legs, and under parts black. The whole insect is rough with greyish and blackish hairs and bristles. Thorax wider than long.

Length $2\frac{1}{2}$ lines.

Hab. Mexico.

CYMATODERA PULCHELLA, n. s.

Yellow. Head keeled down the centre. Thorax with seven warts on the back, six of them placed in two longitudinal parallel lines, the seventh between the lines, and connected, but indistinctly, with the middle wart of each line; the sides of the thorax gibbous on each side. Elytra with a black spot at the base in the middle, from which proceeds a short black line; about the middle a widish black band, narrowed towards the suture, and not reaching the lateral margin. Surface of elytra rather finely and irregularly punctured, and furnished with several hairs. Legs, antennæ, and under side pale yellow.

Length $2\frac{1}{2}$ lines.

Hab. Columbia.

TILLICERA CHALYBEA Westw.

Subcylindrical. Head small. Elytra wider behind, cyaneous, highly polished. Prothorax very gibbous, contracted before the hind margin. Elytra with the basal half striato-punctate, each elytron behind the middle with an elevated smooth purple spot; tips smooth. Antennæ with the joints from the fourth to the tenth dilated, and serrated on the inside, and equal.

Length 5 lines.

Hab. Tenasserim.

TILLICERA ? *MUTILLÆCOLOR*, n. s.

Head and thorax deep pitchy black, closely punctured, when fresh must be covered with ochrey yellow hairs. Elytra at the base ferrugineous red, coarsely pustulato-striated; a band of ochrey yellow hairs across each elytron, sinuated on the posterior margin; this is followed by a large deep black lunule, after which comes a band of ochrey yellow hairs, straight behind and narrowed towards the side of elytra; tip deep black. Metathorax red. Legs black, with numerous greyish bristles. Abdomen and antennæ wanting.

Length $6\frac{1}{4}$ lines.

Hab. N. India.

CLADISCUS GRACILIS, n. s.

Head, thorax, and sides of elytra, from the base to the middle, red. Antennæ, legs, terminal half of elytra, and suture smoky brown. Antennæ branched from the third joint; the branches are directed inwards, and emit a second branch, which is directed downwards; the elytra are very much elongated, and furnished with several punctured striæ commencing at the base, but evanescent considerably before the tip.

Length $4\frac{1}{4}$ Lines.

Hab. N. India. From Captain Boyes's Collection.

CLADISCUS PRINSEPII, n. s.

Head and thorax red. Elytra blackish brown. Abdomen above and below black. Antennæ, tips of femora, tibiæ, and tarsi black; base of femora red. Antennæ branched from the third joint; two branches, considerably longer than the joints, spring from the end of each joint, and, in common with the joints, are delicately ciliated with black hairs. Elytra elongated, and furnished

with several punctured striæ, emitting short hairs; apex of elytra somewhat abrupt. Wings of a uniform black.

Length 5 lines.

Hab. E. Indies.

CLADISCUS LONGIPENNIS Westw.

Elongated, cylindrical, black. Head and prothorax sanguineous. Antennæ black; all the joints from the third strongly serrated within. Prothorax much contracted behind the middle. Elytra testaceous, much elongated, each with ten striæ formed of large contiguous points. Legs black, setose; tarsi 5-jointed, first joint rather larger than the second, the third and fourth dilated beneath.

Length from 5 to $6\frac{1}{2}$ lines.

Hab. India (Himalayas).

TENERUS CINGALENSIS, n. s.

Of a rich vermilion red. Mouth, antennæ, legs, and under side of abdomen black. A spot on the crown of head and the tip of elytra black: shoulders of elytra square; each elytron with at least four longitudinal keels. Head, thorax, and elytra covered with crowded short hairs. Antennæ with the eight terminal joints dilated in an oblong oval form.

Length $7\frac{1}{4}$ lines.

Hab. Ceylon.

TENERUS ABBREVIATUS, n. s.

Elytra not reaching much beyond the middle of abdomen. Forehead, thorax, greater portion of abdomen of a clear, shining, orange red. Elytra, to beyond the middle, of a palish metallic green; end yellow, with a large blue spot in the middle; edged with black. Tip of abdomen, above and below, of a deep metallic blue. Antennæ with a longish process from the end of each joint directed inwards, each process being somewhat thickened towards the end. Legs and under part of metathora of a deep black. Wings blackish.

Length 5 lines.

Hab. Australia.

SERRIGER COFFINI Westw.

Short, wide, very delicately but thickly punctured, sanguineous. Elytra with two blackish blue spots before the middle, the apical half of the same colour. Antennæ black at the base, rufous; the three apical joints, especially the last, widened. Palpi black, all securiform. Legs brownish black; femora rufous at the base. Body beneath black. Abdomen rufous, slightly washed with brown.

Length $6\frac{1}{4}$ lines.

Hab. Mexico.

OMADIUS KAMELIANUS, n. s.

Thorax above somewhat nodulose; two deeply grooved transverse lines, one in front and one behind; the posterior line close to hind margin. Elytra in colour and marking very similar to *O. modestus*. Legs pale yellow; femora of hind leg, when extended, reaching beyond the end of elytra; terminal half of femora of hind legs black, with a yellow spot on upper edge; a black ring round the middle of tibiæ, on front legs nearly obsolete.

Length $5\frac{1}{4}$ lines.

Hab. Philippine Islands. From Mr. Cuming's Collection.

STIGMATIUM GILBERTI, n. s.

Of a dullish ferruginous. The head covered with grey down and bristles, extending even over the eyes. Thorax with a deeply impressed transverse line; the surface very minutely punctured. Elytra with the basal half rather coarsely punctured, the punctures running in lines; the posterior part of elytra smoothish, black; the tip, and an oblong spot before the tip and close to the suture, also covered with greyish hairs slightly bent; there is a somewhat indistinct band of greyish hairs across the elytra, terminating where the black commences.

Length from $3\frac{1}{2}$ to $4\frac{3}{4}$ lines.

Hab. N. Holland, Port Essington.

STIGMATIUM NATALENSE Westw.

Deep pitchy brown. Elytra punctato-striate, the basal part with various spots and patches of greyish hairs between the striæ; a wide, transverse, irregular band of greyish hairs beyond the middle, running down the suture to the tip, and forming a spot there. Thorax scarcely, if at all gibbous on the sides. Face

with numerous long greyish hairs. Under side of abdomen pale ferruginous. Legs pale yellowish brown, with short grey hairs and a few scattered bristles; tarsi of a deeper brown.

Length 5 lines.

Hab. Port Natal. Coll. Westwood.

STIGMATIUM SPECULARE, n. s.

Of a dull brownish colour, more or less covered with scattered hairs and bristles. A largish patch of grey hairs common to both elytra, and situated on the posterior part of elytra. Elytra rather finely striated with punctured lines. Antennæ, palpi, and legs reddish.

Length 3 lines.

Hab. N. Holland (Port Essington).

STIGMATIUM RUFIVENTRE Westw.

Short, thick, black, covered with bristles. Head and prothorax, under a glass, covered with minute punctures; the middle of the head slightly wider. Elytra, at the base towards the costa, of a purplish chestnut, thickly covered with punctures, the punctures arranged in striæ and obliterated before the tip; an irregular band before the middle, and another before the tip, the tip also itself with dull whitish bristles. Antennæ slender, apical joints scarcely thickened. Metasternum and abdomen beneath testaceous.

Length 4 lines.

Hab. Assam.

STIGMATIUM ELAPHROIDES Westw.

Antennæ very long, with brown and narrow yellow rings. Thorax considerably gibbous on the sides, with a deeply impressed transverse line in front; two oblong brown spots on the back of the hind part. Head and thorax with many adpressed silky yellow hairs, and several long blackish bristles. Elytra above covered, for the greater part, with adpressed silky yellow hairs; the apical half with several brown spots, which taken together have the appearance of two twisted brown bands; punctato-striate, with scattered black bristles.

Length 3 lines.

Hab. Ceylon. R. Templeton, Esq.

THANASIMUS SUBSIMILIS, n. s.

Head, thorax, and legs of a reddish yellow. Elytra pale yellow, with two broad, transverse, blackish brown bands; a slight blackish brown line on the edge of the base of the elytra; the posterior band wider than the anterior. Elytra covered with minute punctures; a few slightly raised longitudinal keels down each elytron; tips of elytra at suture brownish black.

Length $4\frac{1}{2}$ lines.

—— ?

THANEROCLERUS ACCINCTUS Newm.

Var. *Clerus viduus* Westw.

Antennæ pitch-coloured. Thorax black. Legs black; femora chestnut at the base.

Length $3\frac{3}{4}$ lines.

Hab. Van Diemen's Land.

Var. *Clerus medianus* Westw.

Somewhat narrow, black, scarcely shining, closely covered with very delicate punctures. Antennæ slightly clavate, short, chestnut, as is the mouth. Labial palpi brown. Prothorax of a blood-red colour. Elytra with a straight white band across the middle, and a small triangular common white spot at the tip. Legs testaceous, with white setæ. Metasternum black. Posterior coxæ testaceous. Posterior margin of segments of abdomen testaceous.

Length $3\frac{1}{4}$ lines.

Hab. Van Diemen's Land.

NOTOXUS CONGRUUS Newm.

Var. *Opilus femoralis* Westw.

Of a pitch-brown. Elytra with striated punctures; a wide transverse yellow band a little behind the middle; tips yellow. Antennæ and legs chestnut; the basal half of the femora pale yellow; antennæ of the male more elongated. Female twice as large as male; antennæ short, and legs duller in colour.

Length from 4 to 6 lines.

Hab. Australia, King George's Sound.

NOTOXUS SUBFASCIATUS Westw.

Somewhat depressed. Head and prothorax pitch-coloured. Antennæ somewhat fulvous, slender; the three terminal joints slightly thicker. Elytra fulvous, shining, striato-punctate; a band before the middle, interrupted at the suture, and another wider one behind the middle of a pitchy black. Legs fulvous; femora duller in colour. Body beneath pitch-coloured. Labial palpi with the last joint very large and of a pale yellow.

Length 4 lines.

Hab. East Indies.

NOTOXUS CASTANEIPENNIS, n. s.

Covered with light ferruginous short hairs. Head, thorax, and abdomen of a blackish brown. Elytra of a uniform pale chestnut-brown. Antennæ yellowish brown. Legs and base of femora pale yellow; tips of femora blackish brown; tibiæ and tarsi reddish brown. Head punctured. Thorax punctured; a transverse grooved line not far from the fore edge, with a short deep grooved longitudinal line on the middle of the back. Elytra with several longitudinal rows of deeply punctured lines.

Length $6\frac{1}{2}$ lines.

Hab. Bengal. Coll. W. W. Saunders.

NOTOXUS APICALIS, n. s.

Of a clear ochraceous yellow. The elytra tipped with black; the black running in an oblique line towards the suture. Femora, from the base to beyond the middle, yellow; tips of femora black; tibiæ and tarsi black. Jaws and antennæ blackish brown. Head and thorax very smooth, shining; an impressed transverse line, sinuated in the middle, close to front edge of thorax. Elytra covered by many closely placed, punctured, parallel lines. Insect covered with short yellowish hairs.

Length $5\frac{1}{2}$ lines.

——? Coll. F. W. Parry.

NOTOXUS UNICOLOR, n. s.

Of a dirty dark brown, densely covered with a silk-like yellowish pubescence. Under side of abdomen with the edges of segments behind margined with yellow. Elytra with rather distant longitudinal lines of punctures not continued to the tip.

Length from $3\frac{3}{4}$ lines to 7 lines.

Hab. India. Mus. E. India House.

NOTOXUS RUFICORNIS Westw.

Elongated, rather narrow. Head and thorax black, with scattered punctures. Antennæ, labrum, and palpi rufo-fulvous. Elytra with the basal half of a bright fulvo-testaceous; the apical half pitch-coloured, with a transverse whitish band behind the middle and interrupted in the middle. Elytra punctato-striate, the tips smooth. Legs black.

Length 4 lines.

NOTOXUS MÆRENS Westw.

Head black, very slightly punctured. Antennæ black, pitchy brown at the base. Palpi moderate. Prothorax black, of the width of the head; behind slightly narrower; the disk in the middle slightly impressed, irregularly punctured. Elytra black, punctato-striate; tips smooth; in the middle a transverse whitish band. Body beneath black. Coxæ of a pitchy chestnut; legs black; tibiæ and tarsi pitchy brown.

Length $2\frac{1}{2}$ lines.

Hab. Australia (Adelaide).

NOTOXUS EPHIPPIGER, n. s.

Of a dingy brownish black; a squarish ochrey yellow spot on the suture, behind which and running into it is a narrow yellow transverse band; from this band, and including it, the surface of the elytra is covered with large punctures arranged in longitudinal lines; end of elytra yellowish, and covered with silky greyish hairs; a tuft of black hairs on the back of each elytron at the base. Antennæ yellow. Legs brownish black; tips of tibiæ and the tarsi yellow.

Length $4\frac{3}{4}$ lines.

Hab. New Holland.

NOTOXUS SEX-NOTATUS Westw.

Elongated, narrow. Legs very long. Head cyaneous, subrugose, with minute punctures; as wide as the prothorax. Antennæ elongated, and with the palpi fulvous. Prothorax cyaneous, shining, slightly rugose; before the middle with an angled impression, and an abbreviated middle dorsal line behind, considerably contracted. Elytra cyaneous at the tip, inclined to purple; the basal half striato-punctate; two oval spots before the middle, an interrupted median band, and two somewhat oblique apical spots, yellowish. Body beneath cyaneous. Legs

black; femora at the base, and tip of the tibiæ and tarsi, fulvous. All the palpi securiform.

Length $3\frac{1}{4}$ lines.

Hab. Van Diemen's Land.

NOTOXUS PULCHER, n. s.

Of a deep metallic purplish blue, with two squarish yellow spots on the back of the elytra before the middle, a yellow band across the middle of elytra not extending to the suture, and an oblique yellow spot on the tip. Antennæ and palpi amber-yellow. Legs amber-yellow. Head thickly punctate. Thorax with scattered punctures; two deeply impressed semicircular lines, one on each side, not touching in the middle; a short deeply impressed longitudinal line down part of the middle of thorax. Elytra with its basal portion with large punctures arranged in lines; the apical half smooth and shining; under side deep black.

Var. *a.* Tips of femora purplish blue.

b. Legs wholly yellow.

Length $4\frac{1}{4}$ lines to $6\frac{1}{2}$ lines.

Hab. Van Diemen's Land. Rev. T. Ewing; R. Butler, Esq.

CLERUS SERICANS Westw.

Somewhat narrow, closely covered with very delicate punctures. Head and prothorax æneous, with bristle like golden hairs. Antennæ black, at the base fulvo-piceous; the last three joints gradually thickened. Elytra of a livid ferruginous, covered with black setæ; a pinkish white much-angled band on the middle (margined in front with black); two subapical spots of a pinkish white. Mouth with the palpi of a tawny yellow. Body beneath of a brassy black. Legs black. Belly of a pitchy red. Tarsi pitchy.

Length $3\frac{1}{2}$ lines.

Hab. Mexico.

CLERUS HOLOSERICEUS, n. s.

Depressed. Elytra somewhat widened behind. Head, thorax, above and below, and legs, black, and more or less covered with black hairs and greyish silky pubescence. Elytra with the basal half of a reddish fawn; that colour on the side and suture extending beyond the middle; a large round black spot at the

base, common to both elytra, between which and the margin on each side is a narrow black spot not reaching to the shoulder; the terminal half of elytra black, with a transverse band near the tip, and the suture at the tip with short grey silky hairs. Abdomen beneath of a dark brownish red.

Length 4 lines.

Hab. California. From the Collection of Mr. Hartweg.

CLERUS LATICINCTUS, n. s.

Of a deep purplish black. Elytra with a wide, clear yellow, transverse band across the middle; the sutural margin black; a roundish yellow spot at the base of each elytron (sometimes, as in Capt. Parry's specimen, wanting). Legs of a yellowish red, the tibiæ more obscure; tarsi black. Head and thorax closely punctured, and thickly covered with short, erect, black hairs. A patch of silky greyish brown hairs at the end of each elytron.

Length from $3\frac{1}{2}$ to 4 lines.

Hab. Brazil.

CLERUS GUTTULUS, n. s.

Elongated, narrow. Head as wide as thorax. Thorax suddenly narrowed behind. Elytra with the sides somewhat sinuated. Mouth, antennæ, and legs yellowish red. Head and thorax black, with greenish reflexions; thickly punctured and hairy. Elytra brown, and striato-punctate at the base, with a short, diagonal, ferruginous band from the shoulder to the suture, and two white, smooth, raised lineolets, one on each side of the band, the outer the longer; a narrow, transverse, white, smooth, raised, linear band behind the middle of the elytra, which are black at the tip, and have green reflexions. Under side of abdomen black; mesosternum and metasternum yellow.

Length $3\frac{1}{2}$ lines.

Hab. Australia.

CLERUS ANTHICOIDES Westw.

Pitchy black. Elytra with four largish yellow spots; two transverse before the middle, and two sub-oblique close to the tip. Antennæ long, yellowish; the four terminal joints gradually thickened, brownish black. Thorax as broad as long, slightly narrowed behind. Legs pitchy brown; tarsi yellow; tibiæ of hind legs with some longish hairs on the outside.

Length $1\frac{1}{2}$ line.

Hab. Ceylon.

TRICHODES DOUGLASIANUS, n. s.

A narrow elongated species. Head, thorax, scutellum, and under parts of body, of a lively metallic green. Elytra of a very dark purple, very much decorated with orange yellow bands, one occupying the shoulder, except on the very angle and on the inner edge about the scutellum; it emits at its lower side a yellow spot, connected with the yellow of the base by a narrow yellow line close to the suture; about the middle of elytra, and connected by a narrow, yellow, lateral, marginal line to the yellow of the base, is a diagonal yellow band which ends towards the suture in a spot which is directed upwards; another diagonal yellow band proceeds from the side of the tip upwards, and is angled towards the suture. Legs of a metallic green, but with many rusty yellow hairs, which give them a ferruginous appearance.

Length $4\frac{1}{4}$ lines to 5 lines.

Hab. California. From Mr. Hartweg's Collection.

TRICHODES HARTWEGIANUS, n. s.

Head, thorax, scutellum, elytra, body, and legs of a dark shining green; the hairs longish and of a rusty grey. The shoulder with a squarish yellow mark, with a green spot on the angle; a little yellow spot towards the suture, not connected with the basal yellow spot, nor with the median, transverse, slightly diagonal, yellow band, not quite touching the suture; a yellow arcuated spot near the tip, the sinus on its hinder margin.

Length $5\frac{1}{4}$ lines.

Hab. California. From Mr. Hartweg's Collection.

AULICUS INSTABILIS Newm.

Var. *Clerus tibialis* Westw.

Elytra at the base and tip more punctured, of a golden green; roughly striato-punctate. Legs of a bluish green; anterior legs within, with the tarsi, fulvous. Antennæ and palpi fulvous. Body beneath of a lively green.

Length 4 lines.

Hab. Australia.

Var. *Clerus castanipes* Westw.

Oblong, depressed, cyaneous, or greenish blue. Head and prothorax, under a glass, covered with punctures. Elytra roughly

DESCRIPTIONS OF NEW SPECIES.

striato-punctate; the base and tip rather smooth; shoulders purple. Legs chestnut, with black bristles; the four posterior tarsi brown. Antennæ short, chestnut; three apical joints dilated. Body beneath of a greenish shining cyaneous.

Length 4 lines.

Hab. Van Diemen's Land.

PHLOIOCOPUS GUERINII, n. s.

Head and thorax of a deep brown, the latter ferruginous in front.

Elytra with the basal half of a pale reddish brown, rather roughly punctato-striate; a narrow, irregular, brown band across the middle, followed by a wider, smooth, yellow band with irregular edges; a still wider, smooth, brown band succeeds this; a narrow yellow sub-apical band, the extreme tip is brown. Antennæ rufous. Legs ferruginous brown; base of femora and soles of tarsi inclined to yellow.

Length 6 lines.

Hab. Africa.

HOPLOCLERUS BIACULEATUS *Westw.*

Somewhat depressed. Elytra wider behind. Head and prothorax smooth. Elytra irregularly covered with punctures; of a lively testaceous; shining, cyaneous; shoulders and tips rufous, the latter at the suture ending in two sharp spines. Antennæ short; the three apical joints wider. Body beneath rufous; the mesosternum and metasternum cyaneous.

Length 3 lines.

Hab. S. Australia.

HYDNOCERA PECTORALIS *Westw.*

Allied to *H. flavo-lineata*, but wider behind and less parallel above, altogether of a pale testaceous; anterior margin of prothorax whitish. Antennæ very short, of a pale testaceous; three last joints short, and wider than the rest. Head, between the eyes, with two impressions. Elytra deeply punctato-striate. Legs yellowish. Thorax black beneath. Abdomen chestnut.

Length $2\frac{1}{2}$ lines.

Hab. Australia.

HYDROCERA FLAVO-VARIA Westw.

Head black, face yellow. Antennæ very short, slightly thickened, yellow. Prothorax with the sides before the middle slightly rounded, of the width of the head, yellow, disk marked with a horse-shoe of a pitchy brown. Elytra elongated, somewhat parallel, subrugose, black; each with an oblong, basal, yellow spot, another somewhat rounded, nearly in the middle; another rounded yellow spot before the tip, and the tip itself yellow. Legs yellow, tips of posterior femora, posterior tibiæ, and posterior tarsi, of a pitchy brown. Body beneath black.

Length $2\frac{1}{2}$ lines.

Hab. Australia (King George's Sound).

HYDROCERA FLAVO-LINEATA Westw.

Small, much elongated. Head and prothorax of a dark steel colour, with scattered punctures; face and labrum whitish. Antennæ shorter than the head, yellow, slightly thickened at the tip. Elytra elongated, somewhat parallel, punctured, disk of each marked with a longitudinal palish yellow streak. Legs somewhat fulvous; femora, especially the posterior, black above; hind tarsi black.

Length 2 lines.

Hab. S. Australia.

HYDROCERA TASMANICA, n. s.

Elongated. Head in front and mouth yellow; top of head black; spot on vertex ferruginous. Thorax depressed above, with a deeply impressed line close to and parallel with the front, smooth, ferruginous. Elytra black; the base pinkish yellow, except on the margin, which is black; a pinkish yellow band across the middle, not continued to the margin; the suture black; tip yellow. Two first pairs of legs yellow; hind legs black; base of femora and tibiæ yellow; under side black. Tip of abdomen black. Antennæ wanting in the specimen.

Length 3 lines.

Hab. Van Diemen's Land.

HYDROCERA PICTA, n. s.

Head smooth, black, with green reflexions. Face immediately above trophi edged with yellow. Antennæ black. Thorax depressed on the back, and suddenly strangulated on the sides behind, with a deep impressed line close to one parallel with the front margin; deeply foveated in the middle; bright yellowish

red. Elytra broader behind than in front, striato-punctate and slightly hairy; greenish metallic blue; base near the shoulder yellowish red; a wide yellowish red band across the elytra, just before the middle, widest on the margin. Prothorax beneath yellowish red; mesothorax and metathorax black. Abdomen beneath at the base red, at the tip black. Legs black, with blue reflexions.

Length $3\frac{1}{4}$ lines.

Hab. Port Philip.

NECROBIA PINGUIS Westw.

Short, thick. Head small, black, closely and densely punctured. Antennæ black; the base and labrum fulvous. Prothorax much wider than the head; the sides in the middle dilated into a rounded tubercle, thickly punctured, some of the punctures larger; rugose, sanguineous. Elytra black, roughly punctured; with a slender nearly straight band in the middle, and another near the tip, whitish. Scutellum rufous. Body beneath, with legs, of reddish testaceous.

Length $3\frac{1}{4}$ lines.

Hab. Van Diemen's Land.

NECROBIA EXIMIA, n. s.

Head and thorax dark shining green, with brassy reflexions. Antennæ and trophi pitchy brown. Elytra with a widish, brilliant purple, transverse band on the middle; the remainder with the brightest combination of green, flame-coloured, and brassy colours passing into each other; under side yellowish green. Legs with a bluish tint; covered with many short black hairs.

Length $2\frac{3}{4}$ lines.

Hab. W. Australia.

OPETIOPALPUS OBESUS Westw.

Small, thick, and convex, somewhat shining, punctured, with black bristles. Head and prothorax of a dull sanguineous. Elytra black, without spots. Antennæ and legs of a reddish chestnut; club of antennæ of a dull colour.

Length $1\frac{1}{2}$ lines.

Hab. E. Indies.

INDEX.

A.

abbreviatus, Ten. 12.
 abdominalis, Cler. 15. 22.
 " Cor. 45.
 " Cyl. 2.
 " Prioc. 5.
 " Than. 15.
 accinctus, Than. 17.
 acerbus, Than. 16.
 acutipennis, Call. 4.
 Adonis, Cler. 16.
 ægra, Hydn. 42.
 æneicollis, Cler. 22.
 æquinoctialis, Phyll. 41.
 affinis, Trich. 27, 28, 29.
 agilis, Till. 2.
 albofasciatus, Till. 2.
 Allelidea, 40.
 alvearius, Trich. 27.
 amabile, Pel. 34.
 ambulans, Till. 2.
 Ammios, Trich. 30.
 amœnum, Pel. 33.
 amœnus, Cler. 25.
 " Syst. 7.
 analis, Ax. 6.
 " Cor. 45.
 angustata, Cym. 9.
 angustatus, Cor. 46.
 " Not. 8, 9.
 " Trich. 29.
 annulatus, Cler. 21.
 annulipes, Cler. 15.
 Anthicides, Pyl. 26.
 anthicoides, Cler. 26.
 antiquus, Cler. 21.
 apiarius, Trich. 28.
 apicale, Pel. 36.
 apicalis, Not. 19.
 " Than. 16.
 apicida, Trich. 28.
 apivorus, Trich. 28.
 Apolopha, 37.
 arachnodes, Cler. 24.
 arcuatus, Cler. 21.
 " Trich. 28.
 arthriticus, Trich. 30.
 artifex, Cler. 20.
 aspera, El. 10.
 assimilis, Cler. 21. 25.
 ater, Cor. 46.

aterrima, Ich. 39.
 atra, Necr. 46.
 attenuata, Hydn. 40.
 Aulicus, 31.
 " Call. 4.
 " Trich. 30.
 auratus, Cler. 31.
 auricollis, Opet. 47.
 auricomus, Till. 6.
 auripenne, Pel. 35.
 auripennis, Op. 35.
 australis, Ol. 19.
 " Zen. 26.
 axillaris, Phyll. 41.
 " Trich. 29.
 Axina, 6.
 azurea, Hydn. 42.
 azureus, Sten. 6.

B.

badium, En. 35.
 balteatus, Cyl. 1.
 " Epi. 38.
 basalis, Cler. 21.
 " Epicl. 10.
 " Hydn. 41.
 " Opil. 32.
 biaculeatus, Hopl. 40.
 bicarinata, Hydn. 40.
 bicinctus, Cler. 23.
 " Pyl. 26.
 bicolor, Call. 4.
 " Necr. 46.
 " Till. 36.
 bifasciatus, Omad. 13.
 " Ten. 12.
 " Trich. 31.
 bifasciellus, Till. 4.
 bilobus, Cler. 20.
 bimaculatus, Cyl. 1.
 " Tæn. 12.
 " Till. 2. 7.
 bipartitus, Not. 32.
 " Till. 3.
 bisignatus, Cler. 24.
 bispinosa, Prioc. 6.
 bitæniata, M. 31.
 bizonatus, Trich. 31.
 bombycinus, Cler. 22.
 Boscii, Cym. 9.
 Brachymorphus, 44.

brachyptera, Hydn. 41.
 brevicollis, Cler. 24.
 brevipennis, Hydn. 42.
 brunneus, Not. 9.
 brunnipes, Cler. 20.
 Buquetii, Cyl. 1.
 " Ep. 38.
 " Not. 18.
 " Pel. 35.
 " Phl. 32.
 " Than. 16.

C.

Cæstron, 24.
 calceata, Ich. 39.
 Callitheres, 4.
 callosa, Eb. 27.
 callosus, Op. 27.
 capensis, Ten. 12.
 " Than. 15.
 carbonarius, Cor. 46.
 " Per. 4.
 Carcelii, Trich. 30.
 carus, Till. 3.
 castanea, Prioc. 5.
 castaneipennis, Not. 19.
 castaneus, Opil. 5.
 castanipes, Cler. 31.
 centromaculatus, Not. 19.
 ceraceus, Trich. 30.
 chalybæus, Cor. 45.
 chalybea, Till. 11.
 Chariessa, 44.
 chilensis, Not. 9.
 chinensis, Not. 15.
 cicindeloides, Stigm. 13.
 cincta, Hydn. 41.
 cinctipennis, Col. 7.
 cinctiventris, Prioc. 5.
 cinctum, En. 33.
 cinereo-pilosus, Cler. 23.
 cingulata, Ten. 12.
 cingulata, Cym. 9.
 cingulatus, Till. 9.
 circumcinctum, En. 34.
 Cladiscus, 11.
 Cleridæ, 1.
 cleroides, Pel. 36.
 Cleronomus, 7.
 Clerus, 20.
 clytoides, Cler. 40.

coarctatus, Pez. 40.
 cœruleus, Not. 45.
 Coffini, Serr. 13.
 collare, Pel. 34.
 collaris, Opet. 47.
 " Till. 3.
 Colombiæ, Cler. 22.
 columbianus, Cler. 21.
 columbicus, Than. 15.
 Colyphus, 7.
 compressicornis, Till. 3.
 comptus, Cler. 20.
 conferta, Hydn. 42.
 conflagrata, Cym. 8.
 conflagratus, Till. 8.
 confusus, Not. 8.
 " Than. 16.
 congruus, Not. 19.
 corallinus, Trich. 28.
 Corynetes, 45.
 crabronarius, Cler. 21.
 crabroniformis, Trich. 28.
 crassipes, Not. 6.
 crassus, Cler. 25.
 cribricollis, Nat. 16.
 cribripenne, Pel. 36.
 cribripennis, Pros. 43.
 " Trich. 28.
 crinipenne, En. 39.
 crinitum, En. 34.
 cruciata, Hydn. 41.
 cruciatus, Cler. 25.
 cruentatus, Cler. 21.
 " Not. 19.
 ctenostomoides, All. 40.
 curtipennis, Hydn. 42.
 curvipennis, Cler. 40.
 cyaneus, Cyl. 1.
 " Phyll. 41.
 cyaneo-maculatum, En. 33.
 cyanopterus, Ten. 12.
 Cyldrus, 1.
 cylindricollis, Cym. 9.
 Cymatodera, 8.

D.

Dahlii, Trich. 27.
 damicornis, Orth. 44.
 Dasytoides, Syst. 7.
 dauci, Trich. 39.
 decorata, Prioc. 5.
 decorum, En. 34.
 decussatus, Cler. 21.
 defunctorum, Nocr. 46.
 Dejeanii, Trich. 30.
 Denops, 2.
 depressus, Cler. 26.
 dentatum, En. 32.
 dentatus, Derm. 32.
 Derestenus, 8.
 Dermestoides, Cler. 17.
 dilatatum, En. 43.
 dimidiatipennis, Ich. 39.
 dimidiatus, Not. 32.
 " Psych. 11.
 " Ten. 12.
 discoidalis, Cym. 9.

discolor, Cor. 45.
 distinctus, Cler. 20.
 " Till. 4.
 " Trich. 29.
 distrophus, Epi. 38.
 divisa, Ich. 39.
 domesticus, Not. 18.
 dorsalis, Not. 18.
 Douglasianus, Trich. 31.
 Dregei, Not. 18.
 dubius, Cler. 24.
 " Than. 14.
 duodecim-maculatum, En. 37.
 duodecim-punctatus, Epi. 37.
 Duponti, Plat. 43.
 " Pyt. 43.

E.

Eburiphora, 27.
 elaphroides, Stigm. 14.
 Eleale, 10.
 elegans, Trich. 28.
 elegantulus, Trich. 29.
 Ellipotoma, 40.
 elongata, Chrys. 2.
 elongatus, Plat. 26.
 " Till. 2.
 Enoplioides, Ich. 39.
 Enophium, 32.
 ephippiger, Not. 19.
 Epiclides, 9.
 Epiphlaeus, 37.
 episcopalis, Aul. 31.
 Erichsonii, Lar. 47.
 Erymanthus, 17.
 erythrocephalus, Cler. 14.
 erythrodera, Leb. 43.
 erythrogaster, Cler. 24.
 erythropus, Cler. 20.
 Eurymetopum, 42.
 Evenus, 39.
 eximia, Nocr. 47.

F.

fasciatum, En. 43.
 fasciatus, Cler. 14.
 " Cyl. 1.
 " Ep. 37.
 " Opil. 27.
 fasciculatum, Pel. 33.
 " Trog. 17.
 fasciculatus, Cler. 17.
 fasciolatus, Than. 16.
 fastigiatus, Till. 6.
 fatuus, Pyl. 26.
 favarius, Trich. 29.
 femoralis, Cler. 14.
 " Opil. 19.
 filiformis, Ev. 39.
 fimbrilatum, P. 36.
 Fischeri, Cler. 24.
 flaveolatum, En. 43.
 flavicornis, Not. 18.

flavo-cinctus, Trich. 30.
 flavo-limbatus, Pel. 33.
 flavo-lineata, Hydn. 42.
 flavo-signatus, Cler. 20.
 flavo-varia, Hydn. 42.
 flavum, Enopl. 34.
 formicarius, Than. 14.
 formicaroides, Attel. 14.
 foveolatus, Xyl. 10.
 frontalis, Not. 19.
 fugax, En. 34.
 fulgens, Zen. 26.
 fulvipes, Eur. 43.
 funebris, Not. 17.
 fuscicornis, Trich. 30.

G.

Gallerucoides, Pel. 35.
 Gambiensis, Cler. 25.
 Gayi, Epic. 9.
 gemmatus, Erym. 17.
 geniculatus, Cor. 45.
 germanus, Not. 19.
 giganteus, Not. 8.
 gigas, Not. 18.
 Gilberti, Stigm. 13.
 Goryi, Plat. 43.
 gracilis, Clad. 11.
 griseo-pilosus, Cler. 20.
 Guerinii, Phl. 32.
 gulo, Trich. 28.
 guttulus, Cler. 25.

H.

Hanetii, Not. 3.
 Hartwegianus, Trich. 31.
 helopioides, Phil. 36.
 hilaris, Till. 3.
 hirtulum, Pel. 36.
 hispanicus, Trich. 29.
 histrio, Cler. 23.
 holosericeus, Cler. 25.
 honestum, Trog. 17.
 honestus, Cler. 17.
 Hopei, Cym. 8.
 Hopfneri, Cler. 21.
 Hoploclerus, 40.
 humerale, Pel. 34.
 humeralis, Epi. 38.
 " Hydn. 41.
 " Trich. 30.
 hyalinus, Till. 2.
 Hydnocera, 40.

I.

ibidioides, Cym. 8.
 Ichnea, 38.
 ichneumoneus, Cler. 22.
 idoneus, Xyl. 10.
 illustris, Trich. 29.
 imperialis, Cler. 23.
 inanis, Lem. 40.
 indicus, Om. 13.

indicus, Than. 15.
 inornata, Cym. 9.
 inornatus, Till. 9.
 insignis, Trich. 29.
 instabilis, Aul. 31.
 interceptus, Col. 7.
 interruptus, Col. 7.
 " Opil. 32.
 " Trich. 28.
 Iodamus, 4.

J.

javana, Till. 11.
 javanus, Not. 13.
 Joannisii, Cler. 4.
 jucundus, Cler. 22.

K.

Kamelianus, Om. 13.
 Kirbyi, Pel. 35.

L.

Lacordairei, Cler. 25.
 læta, Cym. 8.
 lætus, Not. 8.
 " Cler. 23.
 lævigatus, Cler. 20.
 Lafertei, Trich. 29.
 laminatus, Trich. 29.
 Lampyroides, Pel. 33.
 Laplacei, Nat. 16.
 Laportei, Cler. 23.
 Laricobius, 47.
 Lasiodera, 35.
 laterale, En. 39.
 laticinctus, Cler. 25.
 latifasciatus, Trich. 20.
 latus, Than. 16.
 Lebasiella, 43.
 Lebasii, Par. 47.
 Lelieurii, Cler. 32.
 Lemidia, 40.
 lepida, Leb. 43.
 lepidus, Cler. 25. 28.
 Leprieurii, Not. 5.
 Lesueurii, Cler. 21. 32.
 leucaspis, Xyl. 10.
 leucophæum, En. 34.
 leucopsideus, Trich. 30.
 liberatum, En. 36.
 limbata, Hydn. 41.
 lineatocollis, Hydn. 41.
 " Ten. 12.
 lituratum, Pel. 36.
 livida, Hydn. 40.
 longicollis, Cym. 9.
 " Den. 2.
 longipennis, Clad. 11.
 longus, Cler. 24. 31.
 " Till. 6.
 Louvelii, Call. 4.
 luctuosum, Pel. 33.
 luctuosus, Cler. 20.

lunatus, Cler. 22.
 luridus, Opet. 47.
 luscus, Cler. 24.
 lycoides, Ich. 38.
 " Plat. 43.
 lymexylonoides, En. 39.

M.

Macrotelus, 37.
 maculatum, Eur. 42.
 maculicollis, Cler. 25.
 madagascariensis, Cler. 26.
 malthinus, Hydn. 42.
 Marchiæ, Crypt. 2.
 marginatum, En. 33.
 marginella, Leb. 44.
 marginellum, En. 39.
 marginellus, Epi. 38.
 marginicollis, Prioc. 5.
 marginipenne, Pel. 34.
 marmorata, Cym. 8.
 marmoratus, Than. 15.
 " Till. 8.
 medianus, Cler. 17.
 megacephalus, Cyl. 1.
 megasceloides, Cler. 40.
 megatoma, Mon. 37.
 melanurum, En. 38.
 metalloideum, En. 33.
 mexicanus, Cler. 21.
 Miersiana, Prioc. 5.
 Miletus, Not. 8.
 miniaceus, Cler. 22.
 minutus, Cler. 20.
 minutus, Cler. 22.
 mitis, Than. 15.
 modesta, Cym. 8.
 modestus, Cler. 13.
 mœrens, Not. 19.
 mœstus, Cler. 24.
 mollis, Not. 18.
 Monophylla, 37.
 mucoreus, Ep. 37.
 Muisca, 31.
 mumiarum, Necr. 47.
 mutillæcolor, Till. 11.
 mutillarius, Than. 14.
 myops, Cler. 21.
 myrmecodes, Cler. 3.
 mysticus, Cler. 21.

N.

natalense, Stigm. 13.
 Natalis, 16.
 nebulosus, Cler. 20.
 " Omad. 13.
 Necrobia, 46.
 Nero, Aul. 31.
 nigricans, Cler. 20.
 nigricorne, En. 35.
 nigrinus, Cyl. 1.
 nigripenne, En. 39. 44.
 nigripes, Cler. 23.
 nigriventris, Cler. 22.
 nigrocinctus, Cler. 23.

nigro-maculatus, Cler. 25.
 nigro-signatum, Pel. 35.
 nitens, Lem. 40.
 nitidus, Cler. 20.
 niveum, Pel. 34.
 nobilis, Cler. 21.
 " Trich. 29.
 notatus, Till. 3.
 Notostenus, 44.
 Notoxus, 17.
 novem-maculatus, Tillus, 4.
 nubilus, Ep. 37.
 " Than. 15.
 Nuttalli, Trich. 28. 30.

O.

obesa, Zen. 26.
 obesus, Cler. 26.
 " Opet. 47.
 obliquatus, Trich. 29.
 oblique-fasciatus, Cler. 20.
 obscurus, Cler. 10.
 " Not. 18.
 obsoletum, P. 37.
 obsoletus, Than. 15.
 ochropus, Yl. 26.
 octopunctatus, Trich. 27.
 oculatus, Cler. 24.
 Olesterus, 19.
 Olivieri, Trich. 30.
 Omadii, 13.
 omoplatus, Trich. 30.
 onustum, En. 33.
 opacum, En. 39.
 Opetiopalpus, 47.
 ornatum, Pel. 36.
 ornatus, Cler. 23.
 " Ep. 38.
 " Trich. 31.
 Orthopleurus, 44.

P.

Pachyscelis, 28.
 Pallenis, 4.
 pallens, Eur. 43.
 pallicornis, Cor. 45.
 pallidipennis, Cym. 9.
 pallidus, Not. 19.
 pallipes, Cor. 45.
 " Cyl. 1.
 pannonicus, Trich. 28.
 pantherinus, Ep. 37.
 pantomelas, Aul. 31.
 Paratanetus, 47.
 Passerini, Yl. 26.
 patricia, Eb. 27.
 patricius, Op. 27.
 pectinicornis, Clad. 12.
 pectoralis, Cor. 45.
 " Hydn. 42.
 Pelonium, 33.
 Perilypus, 4.
 personatus, Den. 2.
 Pezoporos, 40.

phaleratus, Cler. 21.
 Phedinus, Trich. 29.
 Philocalus, 7.
 Philyra, 36.
 Phloiocopus, 32.
 Phonius, 7.
 Phyllobænus, 37.
 picipennis, Till. 4.
 picta, Hydn. 42.
 pictus, Ten. 12.
 „ Than. 15.
 pilosus, Lamp. 33.
 pinguis, Necr. 46.
 planatus, Plat. 26.
 plano-notatus, Cler. 20.
 Platyclerus, 26.
 Platynoptera, 43.
 Plocamocera, 38.
 pondicherianus, Cler. 16.
 porcata, Nat. 16.
 porcatus, Not. 16.
 præustum, En. 39.
 „ Pel. 35.
 præustus, Tell. 12.
 principale, En. 35.
 Prinsepî, Clad. 11.
 Priocera, 5.
 proluxa, Cym. 8.
 proluxus, Cler. 13.
 „ Till. 8.
 Prosymnus, 43.
 Ptychopterus, 11.
 pubescens, Till. 3.
 „ En. 39.
 pulchella, Cym. 8.
 pulchellum, Pel. 36.
 pulchellus, Cler. 20.
 „ Till. 6.
 pulcher, Cler. 11.
 „ Not. 19.
 punctata, Hydn. 42.
 „ Nat. 26.
 „ Not. 16.
 punctatissimum, Pel. 36.
 punctato-striatum, Enopl. 34.
 punctatus, Par. 47.
 „ Trich. 29.
 purpureus, Cler. 22.
 pusillus, Cler. 21.
 „ Cor. 45.
 pustulata, Prioc. 5.
 Pylus, 26.
 Pyticera, 43.
 Q.
 quadra, Necr. 46.
 quadratocolis, Orth. 44.
 quadriguttatus, Trich. 31.
 quadrilineatus, Der. 8.
 quadrimaculatus, Phyll. 41.
 „ Than. 15.
 quadrinotatus, Cler. 25.
 quadripunctatum, En. 32.
 quadripunctatus, Trich. 31.
 quadripustulatus, Trich. 31.
 quadrisignatum, Pel. 34.

R.

ramicornis, Char. 44.
 regalis, Not. 8.
 Reichei, Ap. 37.
 „ Eb. 27.
 „ Scrob. 11.
 „ Ser. 12.
 „ Xyl. 10.
 roseicollis, En. 39.
 rosmarus, Cler. 23.
 rubricollis, Cor. 47.
 „ Till. 3.
 rubripes, Cler. 20.
 rufescens, Prioc. 5.
 ruficeps, Than. 14.
 ruficollis, Cler. 23.
 „ Lagria, 2.
 „ Necr. 46.
 ruficornis, Cor. 45.
 „ Not. 19.
 rufipennis, Cler. 17.
 „ Col. 7.
 rufipes, Cler. 14.
 „ En. 36.
 „ Hydn. 41.
 „ Necr. 46.
 rufitarsis, Not. 6.
 rufiventris, Stigm. 14.
 rufiventris, Cler. 23.
 rufulus, Cler. 22.
 rufus, Cler. 22.
 rugosa, El. 10.
 rusticus, Cler. 13.
 Ryparus, 43.

S.

sanguineo-signatus, Trich. 29.
 sanguineus, Cler. 17.
 sanguinicollis, Orth. 44.
 „ Till. 11.
 sanguinipennis, Phon. 7.
 sanguinolentus, Cler. 16.
 scabripennis, Cor. 45.
 scalaris, Cler. 40.
 scenicus, Cler. 22.
 scoparium, Pel. 34.
 Scrobiger, 11.
 scrobilatus, Xyl. 10.
 scutellaris, Opet. 47.
 scutellatum, Pel. 34.
 semi-nigrum, P. 35.
 semistriatus, Cor. 45.
 senilis, Trich. 29.
 sericans, Cler. 25.
 sericella, Ploc. 38.
 sericeus, Epi. 38.
 serrata, Hydn. 41.
 serraticorne, En. 32.
 serraticornis, Attel. 3.
 serricollis, Lagria, 47.
 Serriger, 12.
 sexmaculata, Ax. 6.
 sexpunctatus, Till. 5.
 signaticollis, Col. 7.

signaticollis, Ten. 12.
 signatus, Cler. 24.
 Silbermanni, Cler. 25.
 simplex, Cler. 13.
 sipylus, Cler. 30.
 smyrnensis, Trich. 30.
 sobrinus, Cler. 20.
 spectabilis, Cor. 44.
 spectrum, Not. 13.
 speculare, Stigm. 13.
 speculum, Epi. 38.
 sphægeus, Cler. 24.
 Spinolæ, Cler. 36.
 spinosa, Prioc. 5.
 spinosus, Cler. 5.
 splendidus, Cler. 11.
 steniformis, Hydn. 42.
 Stenoclerus, 6.
 Stenocylidrus, 6.
 Stigmatium, 13.
 strangulatus, Clad. 11.
 subænea, Hydn. 42.
 suberosus, Opil. 32.
 subfasciatus, Not. 18, 19.
 „ Trich. 31.
 subsimilis, Than. 15.
 subtrifasciatus, Trich. 28.
 succinctus, Cyl. 1.
 „ Till. 3, 7.
 sumatrensis, Opil. 13.
 suturalis, Pel. 33.
 suturalis, Hydn. 41.
 „ Ich. 39.
 syriacus, Not. 19.
 „ Trich. 30.
 Systemoderes, 7.

T.

tæniatus, Not. 18.
 Tarsostenus, 27.
 tasmanica, Hydn. 42.
 tenella, Hydn. 41.
 Tenerus, 12.
 tenuicollis, Till. 2.
 tenuiformis, Ell. 40.
 terebraus, Not. 8.
 terminalis, Cal. 7.
 terminata, Mon. 37.
 terminatum, P. 36.
 terminatus, Ten. 12.
 „ Till. 37.
 testaceum, Pel. 35.
 Thanasimus, 14.
 Thaneroclerus, 16.
 Theano, 42.
 Thomasii, Ten. 38.
 thoracicum, En. 44.
 thoracicus, Cler. 19, 25.
 „ Cor. 46.
 Thunbergi, Cor. 45.
 tibialis, C. er. 24, 31.
 „ Necr. 46.
 Tillicera, 11.
 Tilloclerus, 6.
 Tillioidea, 3.
 Tilius, 2.
 tomentosus, Epi. 38.

tomentosus, Ryp. 43.
transversalis, Phyll. 37. 41.
" Till. 3. 18.

Trichodes, 27.
tricolor, Call. 4.
" Cler. 23.
" Phl. 32.
" Till. 3.
trifasciatus, Cler. 22, 23. 36.
" Om. 13.
" Trich. 28.
trinotata, Prioc. 5.
tripagiatus, Cler. 22.
tristis, Not. 17.
Trogodendron, 17.
Trogositoides, Cler. 24.
tropicus, Not. 18.

U.

umbellatarum, Trich. 27.
undata, Cym. 8.
undatulus, Than. 15.
unicinctus, Cler. 3.
unicolor, Not. 19.
unifasciatus, Cler. 3.
" Not. 18.
" Till. 2. 27.
" Trich. 28.
univittatus, Tars. 27.

V.

variabile, Pel. 35.
variabilis, Ten. 12.
" Trich. 29.
variegata, Prioc. 5.
variegatum, En. 35.
variegatus, Cler. 20, 21.
" Epi. 38.
varius, Cler. 23.
Vauthieri, Not. 9.
venator, Cler. 21.
venustus, Till. 6.
Verreauxii, Than. 15.
versicolor, Cler. 21.
" Than. 15.
vestita, Ch. 44.
vetustum, Pel. 34.
vicinus, Trich. 29.
viduus, Call. 4.
" Cler. 17. 24.
villosa, Axina, 5.
violacea, Necr. 46.
violaceus, Cor. 45.
virescens, Than. 16.
" Xyl. 10.
viridi-fasciatus, Trich. 30.
viridipenne, En. 35.
viridipennis, Syst. 7.
viridis, Not. 44.
" Xyl. 10.
vittatum, Pel. 36.

vitticeps, Plat. 43.
vulneratus, Cler. 24.
vysnagæ, Trich. 30.

W.

Weberi, Enopl. 44.
Westwoodii, Cym. 9.

X.

Xanthoceros, 11. 17.
Xilotretus, 10.
Xylobius, 6.
Xylotretus, 10.

Y.

Yliotis, 26.


Z.

zebra, Cler. 25.
" Trich. 28.
Zenithicola, 26.
zonatus, Cler. 22.
" Opil. 3.
" Till. 7.


THE END.

" LONDON:

SPOTTISWOODES and SHAW,
New-street-Square.







SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00239232 2

nhent QL577.B82X
pt. 4 Catalogue of coleopterous insects